2024 PRIOR AUTHORIZATION CRITERIA

TABLE OF CONTENTS

Abiraterone Acetate	
Abraxane	155
Actimmune	13
Acyclovir	14
Adcetris	155
Adempas	237
Aimovig	15
Akeega	261
Alecensa	261
Aliqopa	155
Alosetron Hydrochloride	16
Alunbrig	261
Alymsys	155
Alyq	247
Ambrisentan	239
Apokyn	
Apomorphine Hydrochloride	
Aranesp Albumin Free	
Arcalyst	29
Aripiprazole	26
Aripiprazole Odt	26
Armodafinil	
Asenapine Maleate Sl	26
Atovaquone	
Augtyro	261
Auryxia	
Austedo	35
Austedo Xr	
Austedo Xr Titration Kit	35
Avastin	155
Avita	
Avonex	
Avonex Pen	
Ayvakit	261
Balversa	261
Bavencio	201
Belbuca	
Beleodaq	155
Belsomra	
Benlysta IV	
Benlysta SC	

Benztropine Mesylate	
Besponsa	
Besremi	
Betaseron	
Bexarotene Cap	
Bexarotene Gel	
Blenrep	
Blincyto	
Bortezomib	
Bosentan	
Bosulif	
Braftovi	
Brukinsa	
Budesonide	
Budesonide Er	
Buprenorphine	
Bydureon Bcise	
Byetta	
Cabometyx	
Calquence	
Caprelsa	
Carglumic Acid	
Cerezyme	
Chenodal	
Chlorpromazine Hydrochloride	
Chorionic Gonadotropin	
Cinacalcet Hydrochloride	
Cinryze	
Clemastine Fumarate	
Clobazam	
Clorazepate Dipotassium	
Clozapine	
Clozapine Odt	
Cometriq	
Copaxone	
Copiktra	
Corlanor	
Cosentyx	
Cosentyx Sensoready Pen	
Cosentyx Uno Inj	
Cotellic	
Cresemba	
Crysvita	
Cyltezo	
Cyltezo Kit	
Cyltezo Pen	
Cyltezo Pen Kit	
Cyltezo Pen-Cd/Uc/Hs Starter Pack	

	51
Cyproheptadine Hydrochloride	
Cyramza	
Cystadrops	
Cystagon	
Cystaran	108
Dalfampridine Er	110
Danazol	18
Danyelza	
Darzalex	155
Darzalex Faspro	
Dasatinib	261
Daurismo	
Dayvigo	
Deferasirox (Exjade)	
Deferasirox (Jadenu)	
Depo-Testosterone	
Dexmethylphenidate Hydrochloride	
Diazepam	
Diazepam Intensol	43
Diclofenac Sodium 3% Gel	
Dicyclomine Hydrochloride	
Dihydroergotamine Mesylate Spray	
Dimethyl Fumarate	184
Dimethyl Fumarate Starterpack	184
Diphenoxylate Hydrochloride/Atropine Sulfate	150
Doxorubicin Hydrochloride Liposomal	155
Doxorubicin Hydrochloride Liposomal Droxidopa	
<i>,</i> , , , , , , , , , , , , , , , , , ,	112
Droxidopa	112 113
Droxidopa Dupixent	112 113 137
Droxidopa Dupixent Elelyso	112 113 137 165
Droxidopa Dupixent Elelyso Eligard	112 113 137 165 115
Droxidopa Dupixent Elelyso Eligard Emgality	112 113 137 165 115 155
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti	112 113 137 165 115 155 116
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti Emsam	112 113 137 165 115 155 116 53
Droxidopa Dupixent Elelyso Eligard Emgality. Empliciti. Emsam. Enbrel	112 113 137 165 115 155 116 53 53
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti Emsam Enbrel Enbrel Mini	112 113 137 165 115 155 116 53 53 53
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti. Emsam Enbrel Enbrel Enbrel Mini. Enbrel Sureclick	112 113 137 165 115 115 116 53 53 53 118
Droxidopa Dupixent Elelyso Eligard Emgality. Empliciti. Emsam. Enbrel Enbrel Mini. Enbrel Sureclick. Endari	112 113 137 165 115 116 53 53 53 118 155
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti Emsam Enbrel Enbrel Mini Enbrel Sureclick Endari Enhertu	112 113 137 165 115 115 116 53 53 53 118 155 119
Droxidopa Dupixent Elelyso Eligard Emgality Emgliciti Emsam Enbrel Enbrel Mini Enbrel Sureclick Enhert Enhert Enhert Enhert	112 113 137 165 115 115 116 53 53 53 53 118 155 119 121
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti Emsam Enbrel Mini Enbrel Sureclick Enhertu Enhertu Epclusa Epidiolex	112 113 137 165 115 116 53 53 53 53 118 155 119 121 155
Droxidopa Dupixent Elelyso Eligard Emgality Emgliciti Emsam Enbrel Enbrel Mini Enbrel Sureclick Endari Enhertu Epclusa Epclusa Erbitux	112 113 137 165 115 115 116 53 53 53 118 155 119 121 155 155
Droxidopa	112 113 137 165 115 155 116 53 53 53 53 118 155 119 121 155 155 261
Droxidopa	112 113 137 165 115 115 116 53 53 53 53 118 155 119 121 155 155 261 261
Droxidopa Dupixent	112 113 137 165 116 155 116 53 53 53 53 118 155 119 121 155 155 261 261 261
Droxidopa Dupixent Elelyso Eligard Emgality Empliciti Emsam Enbrel Mini Enbrel Mini Enbrel Sureclick Endari Enhertu Epclusa Epclusa Epidiolex Eribitux Eribulin Eribulin Erivedge Erivedge Erleada Erlotinib Hydrochloride	112 113 137 165 115 115 155 116 53 53 53 118 155 119 155 155 261 261 261

Eysuvis	127
Fanapt	26
Fanapt Titration Pack	26
Farydak	261
Fasenra	128
Fasenra Pen	128
Fentanyl	211
Fentanyl Citrate Oral Transmucosal	129
Fingolimod	185
Fintepla	130
Fluphenazine Decanoate	26
Fluphenazine Hydrochloride	26
Folotyn	155
Forteo	284
Fotivda	261
Fruzaqla	261
Fulphila	97
Fulvestrant	155
Gammagard Liquid	132
Gammagard S/D Iga Less Than 1Mcg/MI	132
Gammaplex	87
Gamunex-C	132
Gattex	134
Gavreto	261
Gazyva	155
Gefitinib	261
Gilotrif	261
Glatiramer Acetate	186
Glatopa	186
Glydo	166
Granix	98
Hadlima	55
Hadlima Pushtouch Pen	55
Haegarda	145
Halaven	155
Haloperidol	26
Haloperidol Decanoate	26
Haloperidol Lactate	26
Harvoni	149
Herceptin	155
Herceptin Hylecta	155
Herzuma	155
Humira	57
Humira Pediatric Crohns Disease Starter Pack	57
Humira Pen	57
Humira Pen-Cd/Uc/Hs Starter	
Humira Pen-Pediatric Uc Starter Pack	57
Humira Pen-Ps/Uv Starter	57

Hydrocodone Bitartrate Er	213
Hydroxyzine Hydrochloride	150
Hydroxyzine Pamoate	150
Ibrance	261
Icatibant Acetate	147
Iclusig	261
Idhifa	261
Ilaris	151
Imatinib Mesylate	261
Imbruvica	261
Imfinzi	202
Imiquimod	153
Inbrija	154
Inlyta	261
Inqovi	261
Inrebic	261
Ivabradine	103
Ivermectin Cream	160
Ivermectin Tablet	161
Iwilfin	261
Jakafi	261
Javygtor	260
Jaypirca	
Jemperli	
Jevtana	
Kadcyla	155
Kalydeco	162
, Kanjinti	
Kesimpta	
Keytruda	
Kineret	
Kisqali	
Kisqali Femara 200 Dose	
Kisqali Femara 400 Dose	
Kisqali Femara 600 Dose	
Korlym	
Koselugo	
Krazati	
Kyprolis	
Lapatinib Ditosylate	
Lazcluze	
Ledipasvir/Sofosbuvir	
Lenalidomide	
Lenvima 10 Mg Daily Dose	
Lenvina 12 Mg Daily Dose	
Lenvina 12 Mg Daily Dose	
Lenvina 14 Mg Daily Dose	
remaind to Mig Duily Dose	202

Lenvima 20 Mg Daily Dose	262
Lenvima 24 Mg Daily Dose	262
Lenvima 4 Mg Daily Dose	262
Lenvima 8 Mg Daily Dose	262
Leukine	
Leuprolide Acetate	165
L-glutamine	118
Libtayo	205
Lidocaine Hcl Gel	
Lidocaine Hcl Jelly	166
Lidocaine Ointment	
Lidocaine Patch	168
Lidocaine Solution	169
Lidocaine/Prilocaine	170
Lidocan	168
Lidocan III	168
Linezolid	
Lonsurf	262
Lorazepam	44
Lorazepam Intensol	44
Lorbrena	262
Loxapine	26
Lumakras	262
Lumoxiti	
Lumryz	174
Lumryz Pak	
Lupron Depot (1-Month)	165
Lupron Depot (3-Month)	
Lupron Depot (4-Month)	165
Lupron Depot (6-Month)	
Lupron Depot-Ped (1-Month)	
Lupron Depot-Ped (3-Month)	
Lupron Depot-Ped (6-Month)	
Lurasidone	26
Lybalvi	26
Lynparza	262
Lytgobi	262
Margenza	155
Matulane	262
Mayzent	188
Mayzent Starter Pack	188
Mekinist	262
Mektovi	262
Memantine Hcl Titration Pak	176
Memantine Hydrochloride	176
Memantine Hydrochloride Er	
Methscopolamine Bromide	150
Methylphenidate Hydrochloride (Methylin)	177

Methylphenidate Hydrochloride (Ritalin)	
Methylphenidate Hydrochloride Er Tablet	178
Methyltestosterone	23
Mifepristone	164
Miglustat	281
Modafinil	
Molindone Hydrochloride	26
Monjuvi	155
Morphine Sulfate Er	215
Mounjaro	181
Mvasi	155
Myalept	193
Mylotarg	156
Natpara	194
Nelarabine	156
Nerlynx	262
Ninlaro	262
Nivestym	100
Noxafil Pak	230
Nubeqa	262
Nuedexta	195
Nuplazid	196
Nurtec	197
Ocaliva	198
Octreotide Acetate	274
Odomzo	262
Ofev	200
Ogivri	156
Ogsiveo	
Ojemda	
Ojjaara	
Ölanzapine	
Olanzapine Odt	
Omnitrope	
Onivyde	
Ontruzant	
Onureg	
Opdivo	
Opsumit	
Oralair	
Orencia	
Orencia Clickject	
Orgovyx	
Orkambi	
Orserdu	
Ofezla	
Otazepam	
Oxycontin	
Олусонын	∠⊥/

Ozempic	
Padcev	156
Paliperidone Er	26
Palynziq	226
Panretin	
Pazopanib	
Pegasys	228
Pemazyre	262
Pemetrexed	156
Perjeta	
Perphenazine	26
Phesgo	156
Pimecrolimus	31
Pimozide	
Piqray 200Mg Daily Dose	262
Piqray 250Mg Daily Dose	262
Piqray 300Mg Daily Dose	262
Pirfenidone	229
Plegridy	189
Plegridy Starter Pack	189
Polivy	156
Pomalyst	262
Portrazza	156
Posaconazole Dr	230
Posaconazole Inj	230
Posaconazole Susp	230
Poteligeo	156
Pregnyl	95
Procrit	
Prolastin-C	17
Prolia	
Promacta	235
Promethazine Hcl Plain	
Promethazine Hydrochloride	
Promethegan	
Pyrimethamine	
Qinlock	
Quetiapine Fumarate	
Quetiapine Fumarate Er	
Quinine Sulfate	
Regranex	
Renflexis	
Repatha	
Repatha Pushtronex System	
Repatha Sureclick	
Retacrit	
Retevmo	
Revlimid	
	205

Rexulti	
Rezlidhia	263
Riabni	
Rinvoq	
Rinvoq Lq	
Risperidone	
Risperidone Odt	26
Rituxan	70
Rituxan Hycela	72
Roflumilast	257
Romidepsin	156
Rozlytrek	263
Rubraca	263
Ruxience	74
Rybelsus	258
Rybrevant	156
Rydapt	263
Sajazir	147
Sapropterin Dihydrochloride	260
Sarclisa	
Scemblix	263
Scopolamine	
Secuado	
Signifor	
Signifor Lar	
Sildenafil	
Sivextro	
Skyrizi	
Skyrizi Pen	
Sodium Oxybate	
Sodium Phenylbutyrate	
Sofosbuvir/Velpatasvir	
Somatuline Depot	
Somavert	
Sorafenib	
Sovaldi	
Spravato 56Mg Dose	
Spravato 84Mg Dose	
Sprycel	
Stelara	
Stelara	
Strensig	
Sunitinib Malate	
Sympazan	
Synribo	
Tabrecta	
Tacrolimus	
Tadalafil 20Mg Tablet	247

Tafinlar	263
Tagrisso	263
Talzenna	263
Tasigna	263
Tasimelteon	283
Tazarotene	289
Tazorac	289
Tazverik	263
Tecentriq	207
Tepmetko	263
Teriparatide	284
Testosterone	24
Testosterone Cypionate	19
Testosterone Enanthate	21
Testosterone Pump	24
Testosterone Topical Solution	24
Tetrabenazine	286
Thalomid	263
Thioridazine Hcl	26
Thiothixene	26
Tibsovo	263
Tobramycin Neb	287
, Tolvaptan	
Torpenz	
Tracleer	
Tramadol Hcl Er	219
Trazimera	156
Trelstar Mixject	
Tremfya	
Tretinoin 10Mg Cap	
Tretinoin Cream, Gel	
Tridacaine	
Trientine Hydrochloride	
Trifluoperazine Hcl	
Trikafta	
Trodelvy	
Trulicity	
Truqap	
Truxima	
Tukysa	
Turalio	
Tymlos	
Tysabri	
Ubrelvy	
Udenyca	
Udenyca Onbody	
Unituxin	
Vanflyta	
vannyta	205

Vectibix	. 156
Vegzelma	. 156
Venclexta	
Venclexta Starting Pack	. 263
Ventavis	. 249
Versacloz	26
Verzenio	. 263
Viberzi	. 299
Victoza	300
Vitrakvi	263
Vizimpro	263
Vonjo	263
Voranigo	263
Voriconazole	301
Vosevi	303
Votrient	263
Vpriv	139
Vumerity	192
yndamax	
Vyndagel	
Vyxeos	
Welireg	
Xalkori	
Xeljanz Solution	
Xeljanz Tablet	
Xeljanz Xr	
Xermelo	
Xgeva	
Xifaxan	
Xolair	
Xospata	
Xosputa	
Xpovio 100 Mg Once Weekly	
Xpovio 100 Mg Once Weekly	
Xpovio 40 Mg Twice Weekly	
Xpovio 60 Mg Once Weekly	
Xpovio 60 Mg Twice Weekly	
Xpovio 80 Mg Once Weekly	
Xpovio 80 Mg Twice Weekly	
Xtandi	
Yargesa	
Yervoy	
Yondelis	
Yonsa	
Zaltrap	
Zejula	
Zelboraf	
Zepatier	313

Zepzelca	156
Ziextenzo	
Ziprasidone Mesylate	
Zirabev	
Zokinvy	
Zolinza	
Ztlido	
Zydelig	
Żykadia	
Żynlonta	
Żyprexa Relprevv	

Prior Authorization Group Description: Actimmune PA Drug Name(s) Actimmune Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require BOTH of the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Prior Authorization Group Description: Acyclovir Topical PA Drug Name(s) Acyclovir Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Prior Authorization Group Description: Aimovig PA Drug Name(s) Aimovig Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of migraine AND 2. The requested agent is being used for migraine prophylaxis AND 3. Patient has 4 migraine headaches or more per month AND 4. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis Criteria for renewal approval require ALL of the following: 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND 2. Patient has a diagnosis of migraine AND 3. The requested agent is being used for migraine prophylaxis AND 4. Patient has had clinical benefit with the requested agent AND 5. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Alosetron PA

Drug Name(s)

Alosetron Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of irritable bowel syndrome with severe diarrhea (IBS-D) AND
- 2. Patient's sex is female AND
- 3. Patient exhibits at least ONE of the following:
 - a. Frequent and severe abdominal pain/discomfort OR
 - b. Frequent bowel urgency or fecal incontinence OR
 - c. Disability or restriction of daily activities due to IBS AND

4. Prescriber has ruled out anatomic or biochemical abnormalities of the gastrointestinal tract

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Alpha-1-Proteinase Inhibitor PA - Prolastin-C

Drug Name(s)

Prolastin-C Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information: Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema

AND 2. Patient has a pre-treatment serum alpha-1 antitrypsin (AAT) level less than 11 micromol/L (80 mg/dL

by immunodiffusion or 57 mg/dL using nephelometry) AND

3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema AND

3. Patient has had clinical benefit with the requested agent AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Anabolic Steroid PA – Danazol

Drug Name(s)

Danazol

Indications: All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:

A. Patient has an FDA labeled indication for the requested agent OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Updated 10/2024

Androgen Injectable PA - testosterone cypionate

Drug Name(s)

Depo-Testosterone

Testosterone Cypionate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient's sex is male with AIDS/HIV-associated wasting syndrome AND BOTH of the following:
 - i. ONE of the following:

a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR

b. Body mass index less than 20 kg/m2 OR

c. At least 5% total body cell mass (BCM) loss within 6 months OR

- d. BCM less than 35% of total body weight and BMI less than 27 kg/m2 AND
- ii. All other causes of weight loss have been ruled out OR
- B. Patient's sex is female with metastatic/inoperable breast cancer OR
- C. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism OR
- D. Patient's sex is male and is an adolescent with delayed puberty AND
- 2. If the patient's sex is a male, ONE of the following:

A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:

i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR

ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR

B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:

i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR

ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND

3. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent **Age Restriction**:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications **Other Criteria**:

Androgen Injectable PA - testosterone enanthate

Drug Name(s)

Testosterone Enanthate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient's sex is male with AIDS/HIV-associated wasting syndrome AND BOTH of the following: i. ONE of the following:
 - a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR
 - b. Body mass index less than 20 kg/m2 OR
 - c. At least 5% total body cell mass (BCM) loss within 6 months OR
 - d. BCM less than 35% of total body weight and BMI less than 27 kg/m2 AND
 - ii. All other causes of weight loss have been ruled out OR
 - B. Patient's sex is female with metastatic/inoperable breast cancer OR
 - C. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism OR
 - D. Patient's sex is male and is an adolescent with delayed puberty AND
- 2. If the patient's sex is a male, ONE of the following:
 - A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:
 - i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR
 - ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR
 - B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:
 - i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR
 - ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND
- 3. ONE of the following:
 - A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent **Age Restriction:**

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications

Androgen Oral PA

Drug Name(s)

Methyltestosterone

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Patient's sex is male with cryptorchidism OR
 - B. Patient's sex is male with hypogonadism OR
 - C. Patient's sex is male and is an adolescent with delayed puberty OR
 - D. Patient's sex is female with metastatic/inoperable breast cancer AND
- 2. If the patient's sex is male, ONE of the following:

A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:

i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR

ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR

B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:

i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR

ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND

3. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent **Age Restriction:**

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications **Other Criteria:**

Androgen Topical PA

Drug Name(s)

Testosterone

Testosterone Pump

Testosterone Topical Solution

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

A. Patient has AIDS/HIV-associated wasting syndrome AND BOTH of the following:

i. ONE of the following:

a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR

b. Body mass index less than 20 kg/m2 OR

c. At least 5% total body cell mass (BCM) loss within 6 months OR

d. In men: BCM less than 35% of total body weight and BMI less than 27 kg/m2 $\ensuremath{\mathsf{OR}}$

e. In women: BCM less than 23% of total body weight and BMI less than 27 kg/m2 AND

ii. All other causes of weight loss have been ruled out OR

B. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism AND 2. If the patient's sex is male, ONE of the following:

A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:

i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR

ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR

B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:

i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR

ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND

3. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has submitted information in support of therapy with more than one agent **Age Restriction**:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria: **Prior Authorization Group Description:** Antipsychotics PA Drug Name(s) Aripiprazole Aripiprazole Odt Asenapine Maleate Sl Chlorpromazine Hydrochloride Clozapine Clozapine Odt Fanapt Fanapt Titration Pack Fluphenazine Decanoate Fluphenazine Hydrochloride Haloperidol Haloperidol Decanoate Haloperidol Lactate Loxapine Lurasidone Lybalvi Molindone Hydrochloride Olanzapine Olanzapine Odt Paliperidone Er Perphenazine Pimozide Quetiapine Fumarate Quetiapine Fumarate Er Rexulti Risperidone **Risperidone Odt** Secuado Thioridazine Hcl Thiothixene Trifluoperazine Hcl Versacloz Ziprasidone Mesylate Zyprexa Relprevv Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** PA does NOT apply to patients less than 65 years of age. Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent OR c. ONE of the following:

i. Patient has a diagnosis other than dementia-related psychosis or dementia related behavioral symptoms OR

ii. Patient has dementia-related psychosis or dementia related behavioral symptoms AND BOTH of the following:

1. Dementia-related psychosis is determined to be severe or the associated behavior puts the patient or others in danger AND

2. Prescriber has documented that s/he has discussed the risk of increased mortality with the patient and/or the patient's surrogate decision maker

Approval authorizations will apply to the requested medication only.

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Apomorphine Inj PA

Drug Name(s)

Apokyn

Apomorphine Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. The requested agent will be used to treat acute, intermittent hypomobility, "off" episodes ("end of dose wearing off" and unpredictable "on/off" episodes) associated with advanced Parkinson's disease AND

2. The requested agent will be used in combination with agents used for therapy in Parkinson's disease (e.g., levodopa, dopamine agonist, monoamine oxidase B inhibitor) AND

3. Patient will NOT be using the requested agent in combination with a 5-HT3 antagonist (e.g.,

ondansetron, granisetron, dolasetron, palonosetron, alosetron)

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Arcalyst PA

Drug Name(s)

Arcalyst

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. ONE of the following:

A. Patient has been diagnosed with Cryopyrin-Associated Periodic Syndromes (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR B. BOTH of the following:

i. Patient has a diagnosis of deficiency of interleukin-1 receptor antagonist AND

ii. The requested agent is being used for maintenance of remission OR

C. BOTH of the following:

i. Patient has a diagnosis of recurrent pericarditis AND

ii. The requested agent is being used to reduce the risk of recurrence AND

2. Patient will NOT be using the requested agent in combination with another biologic agent

Age Restriction:

For diagnosis of CAPS including FCAS or MWS, patient is 12 years of age or over

For diagnosis of recurrent pericarditis and reduction in risk of recurrence, patient is 12 years of age or over

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Prior Authorization Group Description: Armodafinil PA Drug Name(s) Armodafinil Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require BOTH of the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another target agent (i.e., modafinil) Age Restriction: Patient is 17 years of age or over **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Atopic Dermatitis PA – Pimecrolimus

Drug Name(s)

Pimecrolimus

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

- 1. Patient has a diagnosis of atopic dermatitis or vulvar lichen sclerosus AND ONE of the following:
 - A. Patient has tried and had an inadequate response to a topical corticosteroid or topical corticosteroid combination preparation (e.g., hydrocortisone, triamcinolone) OR
 - B. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR

C. Patient has an FDA labeled contraindication to a topical corticosteroid or topical corticosteroid combination preparation OR

2. Patient has a diagnosis of facial seborrheic dermatitis associated with HIV infection AND BOTH of the following:

A. Patient is currently on an antiretroviral treatment regimen AND

B. ONE of the following:

i. Patient has tried and had an inadequate response to a topical corticosteroid or topical antifungal treatment (e.g., hydrocortisone, triamcinolone, ketoconazole) OR

ii. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical antifungal treatment OR

iii. Patient has an FDA labeled contraindication to a topical corticosteroid or topical antifungal treatment OR

3. Patient has an indication that is supported in CMS approved compendia for the requested agent **Age Restriction:**

- Prescriber Restrictions:
- **Coverage Duration:**

Approval will be for 12 months **Other Criteria**:

Atopic Dermatitis PA – Tacrolimus

Drug Name(s)

Tacrolimus Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require ONE of the following: 1. Patient has a diagnosis of atopic dermatitis AND ONE of the following: A. Patient has tried and had an inadequate response to a topical corticosteroid or topical corticosteroid combination preparation (e.g., hydrocortisone, triamcinolone) OR B. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR C. Patient has an FDA labeled contraindication to a topical corticosteroid or topical corticosteroid combination preparation OR 2. Patient has an indication that is supported in CMS approved compendia for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:**

Approval will be for 12 months Other Criteria:

Atovaquone PA

Drug Name(s)

Atovaquone

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. BOTH of the following:
 - i. ONE of the following:
 - 1. Patient has a diagnosis of mild-to-moderate Pneumocystis jirovecii pneumonia OR
 - 2. Patient is using the requested agent for prevention of Pneumocystis jirovecii pneumonia AND
 - ii. ONE of the following:
 - 1. Patient has an intolerance or hypersensitivity to
 - trimethoprim/sulfamethoxazole (TMP/SMX) OR
 - 2. Patient has an FDA labeled contraindication to
 - trimethoprim/sulfamethoxazole (TMP/SMX) OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested

agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Auryxia PA

Drug Name(s)

Auryxia Indications: All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used as iron replacement therapy to treat iron deficiency anemia AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has a diagnosis of hyperphosphatemia AND BOTH of the following:

A. Patient has chronic kidney disease AND

B. Patient is on dialysis

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Austedo PA

Drug Name(s)

Austedo

Austedo Xr

Austedo Xr Titration Kit

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of chorea associated with Huntington's disease AND BOTH of the following:

- i. ONE of the following:
 - 1. Patient does NOT have a current diagnosis of depression OR
 - 2. Patient has a current diagnosis of depression and is being treated for depression AND
- ii. ONE of the following:
 - 1. Patient does NOT have a diagnosis of passive suicidal ideation OR

2. Patient has a diagnosis of passive suicidal ideation and must NOT be actively suicidal OR

B. Patient has a diagnosis of tardive dyskinesia AND ONE of the following:

i. Prescriber has reduced the dose of or discontinued any medications known to cause tardive dyskinesia (i.e., dopamine receptor blocking agents) OR

ii. Prescriber has provided clinical rationale indicating that a reduced dose or

discontinuation of any medications known to cause tardive dyskinesia is not appropriate AND

2. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) AND

3. Patient will NOT be using the requested agent in combination with reserpine

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Belsomra PA Drug Name(s) Belsomra Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Benlysta IV PA

Drug Name(s)

Benlysta IV

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. ONE of the following:

a. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:

i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR

b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:

i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND

- 2. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. ONE of the following:
 - a. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:

i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND

- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

For diagnosis of active systemic lupus erythematosus (SLE) disease, patient is 5 years of age or over. For diagnosis of active lupus nephritis (LN), patient is 5 years of age or over.

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benlysta SC PA

Drug Name(s)

Benlysta SC

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. ONE of the following:

a. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:

i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR

b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:

i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND

- 2. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. ONE of the following:
 - a. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:

 Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:

i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND

- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another biologic agent AND
- 5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

For diagnosis of active systemic lupus erythematosus (SLE) disease, patient is 5 years of age or over. For diagnosis of active lupus nephritis (LN), patient is 18 years of age or over.

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Clobazam

Drug Name(s)

Clobazam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

- a. Seizure disorder OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Clorazepate

Drug Name(s)

Clorazepate Dipotassium

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

- a. Seizure disorder OR
- b. Anxiety disorder AND ONE of the following:

1) Patient has tried and has an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin

norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR

3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR

- c. Alcohol withdrawal OR
- d. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Diazepam

Drug Name(s)

Diazepam

Diazepam Intensol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Anxiety disorder AND ONE of the following:

1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin

norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR

3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR

c. Skeletal muscle spasms OR

d. Alcohol withdrawal OR

e. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Lorazepam

Drug Name(s)

Lorazepam

Lorazepam Intensol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Anxiety disorder AND ONE of the following:

1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin

norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR

3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Oxazepam

Drug Name(s)

Oxazepam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. BOTH of the following:
 - i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Anxiety disorder AND ONE of the following:

1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin

norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR

3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR

b. Alcohol withdrawal OR

c. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Benzodiazepines PA – Sympazan

Drug Name(s)

Sympazan

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Bexarotene Gel PA

Drug Name(s)

Bexarotene Gel

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

- C. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:

a. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions AND

b. ONE of the following:

i. Patient has refractory or persistent disease despite a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR

ii. Patient has an intolerance or hypersensitivity to a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiguimod) OR

iii. Patient has an FDA labeled contraindication to a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR

2. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., dermatologist, oncologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND

iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Patient has had clinical benefit with the requested agent AND

ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., dermatologist, oncologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND

iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Biologic Immunomodulators PA – Cosentyx

Drug Name(s)

Cosentyx

Cosentyx Sensoready Pen

Cosentyx Uno Inj

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria: Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis or plaque psoriasis

NO prerequisites are required for diagnoses of ankylosing spondylitis, enthesitis related arthritis, hidradenitis suppurativa, or non-radiographic axial spondyloarthritis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Biologic Immunomodulators PA – Cyltezo

Drug Name(s)

Cyltezo

Cyltezo Kit

Cyltezo Pen

Cyltezo Pen Kit

Cyltezo Pen-Cd/Uc/Hs Starter Pack

Cyltezo Pen-Ps Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR

C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others **Other Criteria:**

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA – Enbrel

Drug Name(s)

Enbrel

Enbrel Mini

Enbrel Sureclick

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria: Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile psoriatic arthritis, or juvenile idiopathic arthritis

NO prerequisites are required for a diagnoses of ankylosing spondylitis or severe juvenile psoriatic arthritis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, juvenile psoriatic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Biologic Immunomodulators PA – Hadlima

Drug Name(s)

Hadlima

Hadlima Pushtouch Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others **Other Criteria:**

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA – Humira

Drug Name(s)

Humira

Humira Pediatric Crohns Disease Starter Pack

Humira Pen

Humira Pen-Cd/Uc/Hs Starter

Humira Pen-Pediatric Uc Starter Pack

Humira Pen-Ps/Uv Starter

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the

same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others **Other Criteria:**

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA – Kineret

Drug Name(s)

Kineret

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

i. Patient's diagnosis is indicated for preferred biologic immunomodulator agent(s) AND ONE of the following:

a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR

c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria: Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq tablets, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

NO preferred agent is required for diagnoses of Neonatal-Onset Multisystem Inflammatory Disease (NOMID) or Deficiency of Interleukin-1 Receptor Antagonist (DIRA)

Biologic Immunomodulators PA – Orencia

Drug Name(s)

Orencia

Orencia Clickject

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

i. Patient's diagnosis is indicated for preferred biologic immunomodulator agent(s) AND ONE of the following:

a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

- b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR
- c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq, Xeljanz tablets, or Xeljanz oral solution) is required for diagnosis of juvenile idiopathic arthritis

Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq tablets, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

For patients 18 years of age or over, use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Hadlima, Humira, Otezla, Rinvoq, Skyrizi, Stelara, Tremfya, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of psoriatic arthritis

For patients between 6 and less than 18 years of age, use of TWO preferred agents (Cosentyx and Rinvoq) is required for diagnosis of psoriatic arthritis

For patients between 2 and less than 6 years of age, use of ONE preferred agent (Rinvoq) is required for diagnosis of psoriatic arthritis

NO preferred agent is required for diagnosis of prophylaxis of acute graft vs host disease

Biologic Immunomodulators PA – Renflexis

Drug Name(s)

Renflexis

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

i. Patient's diagnosis is indicated for preferred biologic immunomodulator agent(s) AND ONE of the following:

a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR

c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria: Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnosis of rheumatoid arthritis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Hadlima, Humira, Otezla, Skyrizi, Stelara, or Tremfya) is required for diagnosis of psoriatic arthritis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Hadlima, Humira, Otezla, Skyrizi, Stelara, or Tremfya) is required for diagnosis of plaque psoriasis

Use of TWO preferred agents (Cosentyx, Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnosis of ankylosing spondylitis

Use of TWO preferred agents (Cyltezo, Hadlima, Humira, Skyrizi, or Stelara) is required for diagnosis of adult Crohn's disease

Use of TWO preferred agents (Cyltezo, Hadlima, Humira, Skyrizi, Stelara, or Tremfya) is required for diagnosis of adult ulcerative colitis

Use of ONE preferred agent (Cyltezo, Hadlima, or Humira) is required for diagnosis of pediatric Crohn's disease

Only the preferred agent Humira is required for diagnosis of pediatric ulcerative colitis

NO preferred agent is required for diagnosis of adult fistulizing Crohn's disease

Biologic Immunomodulators PA – Riabni

Drug Name(s)

Riabni

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR 2. ALL of the following:

A. ONE of the following:

i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:

 a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR

c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND

B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND

C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq tablets, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA – Rinvoq

Drug Name(s)

Rinvoq

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

i. BOTH of the following:

a. Patient has an FDA labeled indication other than moderate to severe atopic dermatitis for the requested agent AND

b. ONE of the following:

1. Patient's medication history indicates use of preferred TNF agent(s) OR

2. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR

3. Patient has an FDA labeled contraindication to preferred TNF agent(s) OR

4. The request is for an FDA labeled indication that is not covered by preferred TNF agent(s) OR

ii. Patient has a diagnosis of moderate to severe atopic dermatitis AND ONE of the following:

1. Patient's medication history indicates use of TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication OR

2. Patient has an intolerance or hypersensitivity to TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication OR 3. Patient has an FDA labeled contraindication to TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication AND ONE formulary topical calcineurin inhibitor) for the requested indication AND ONE formulary topical calcineurin inhibitor) for the requested indication AND ONE formulary topical calcineurin inhibitor) for the requested indication AND ONE formulary topical calcineurin inhibitor) for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses of ankylosing spondylitis, rheumatoid arthritis, adult psoriatic arthritis, or juvenile idiopathic arthritis

Use of ONE preferred TNF (Cyltezo, Hadlima, or Humira) is required for diagnoses of ulcerative colitis or Crohn's disease

Use of TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) are required for diagnosis of moderate to severe atopic dermatitis

NO preferred TNF agents are required for diagnosis of pediatric psoriatic arthritis

NO preferred TNF agents are required for diagnosis of non-radiographic Axial Spondyloarthritis

Biologic Immunomodulators PA – Rinvoq Solution

Drug Name(s)

Rinvoq Lq

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

- i. Patient's medication history indicates use of preferred TNF agent(s) OR
- ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
- iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) OR
- iv. The request is for an FDA labeled indication that is not covered by preferred TNF agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses adult psoriatic arthritis or juvenile idiopathic arthritis

NO preferred TNF agents are required for diagnosis of pediatric psoriatic arthritis

Biologic Immunomodulators PA – Rituxan

Drug Name(s)

Rituxan

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR 2. ALL of the following:

A. ONE of the following:

i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:

 a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR

c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND

B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND

C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq tablets, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA - Rituxan Hycela

Drug Name(s)

Rituxan Hycela

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND

ii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

iii. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following

i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator ANDiv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

There are no preferred agents required for Rituxan Hycela

Biologic Immunomodulators PA – Ruxience

Drug Name(s)

Ruxience

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR 2. ALL of the following:

A. ONE of the following:

i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:

 a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR

c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND

B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND

C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq tablets, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA – Skyrizi

Drug Name(s)

Skyrizi

Skyrizi Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Use of ONE conventional prerequisite agent is required for diagnoses of Crohn's disease, plaque psoriasis, psoriatic arthritis, or moderate ulcerative colitis

NO prerequisites are required for diagnosis of severe ulcerative colitis

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA – Stelara

Drug Name(s)

Stelara

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, moderate ulcerative colitis, or Crohn's disease

NO prerequisites are required for diagnosis of severe ulcerative colitis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, mercaptopurine

Biologic Immunomodulators PA – Tremfya

Drug Name(s)

Tremfya Indications: All FDA-Approved Indications. Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed ORC. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR

D. Patient's diagnosis does NOT require a conventional prerequisite agent OR

E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR

F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR

G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic

immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, or moderate ulcerative colitis

NO prerequisites are required for diagnosis of severe ulcerative colitis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Biologic Immunomodulators PA - Truxima

Drug Name(s)

Truxima

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR 2. ALL of the following:

A. ONE of the following:

i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:

 a. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

b. Patient has an intolerance or hypersensitivity to preferred biologic immunomodulator agent(s) OR

c. Patient has an FDA labeled contraindication to preferred biologic immunomodulator agent(s) OR

ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND

B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND

C. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

ii. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

iii. Patient will NOT be using the requested agent in combination with another biologic immunomodulator AND

iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred agents (Cyltezo, Enbrel, Hadlima, Humira, Rinvoq tablets, Xeljanz tablets, or Xeljanz XR) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred agents

Biologic Immunomodulators PA - Xeljanz Solution

Drug Name(s)

Xeljanz Solution

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

- i. Patient's medication history indicates use of preferred TNF agent(s) OR
- ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
- iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnosis of juvenile idiopathic arthritis

Biologic Immunomodulators PA - Xeljanz Tablet

Drug Name(s)

Xeljanz Tablet

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

- i. Patient's medication history indicates use of preferred TNF agent(s) OR
- ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
- iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses of psoriatic arthritis, rheumatoid arthritis, juvenile idiopathic arthritis, or ankylosing spondylitis

Use of ONE preferred TNF (Cyltezo, Hadlima, or Humira) is required for diagnosis of ulcerative colitis

Biologic Immunomodulators PA - Xeljanz XR

Drug Name(s)

Xeljanz Xr

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. ONE of the following:

- i. Patient's medication history indicates use of preferred TNF agent(s) OR
- ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
- iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND

3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses of psoriatic arthritis, rheumatoid arthritis, or ankylosing spondylitis

Use of ONE preferred TNF (Cyltezo, Hadlima, or Humira) is required for diagnosis of ulcerative colitis

Bivigam/Flebogamma/Gammaplex/Octagam/Privigen PA

Drug Name(s)

Gammaplex

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

- 1. Patient has ONE of the following diagnoses:
 - A. Primary immunodeficiency [e.g., congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, Xlinked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency] OR
 - B. B-cell chronic lymphocytic leukemia OR multiple myeloma AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
 - iii. Patient has hypogammaglobulinemia OR
 - C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g.,
 - methylprednisolone), or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g.,
 - methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g.,
 - methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., tumor necrosis factor antagonists (e.g., Humira), DMARDS (e.g., methotrexate)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 6 months for indications in Other Criteria, 12 months for all others **Other Criteria**:

G. Myasthenia gravis (MG) AND ONE of the following:

i. Patient is in acute myasthenic crisis OR

ii. Patient has severe refractory MG (e.g., major functional disability/weakness) AND ONE of the following:

a) Patient has failed ONE immunomodulator therapy (i.e., corticosteroid, pyridostigmine, or azathioprine) OR

b) Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR

H. Multiple sclerosis (MS) AND BOTH of the following:

i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND

ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication to TWO MS agents (e.g., Avonex, Betaseron, Copaxone, dimethyl

fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR I. Acquired von Willebrand hemophilia AND ONE of the following:

i. Patient has failed ONE conventional therapy (e.g., desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, or recombinant factor VIIa) OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

J. Refractory pemphigus vulgaris AND ONE of the following:

i. Patient has failed ONE conventional immunosuppressive therapy (e.g., azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR

2. ONE of the following:

A. Patient has another FDA labeled indication for the requested agent OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 6 months approval duration: Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcus, Toxic epidermal necrolysis and Stevens-Johnson syndrome

Drug is also subject to Part B versus Part D review.

Budesonide Oral ER PA – Entocort Drug Name(s) Budesonide Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Budesonide Oral ER PA – Uceris Drug Name(s) Budesonide Er Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Bydureon PA

Drug Name(s)

Bydureon Bcise

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

- 2. Patient had an ineffective treatment response to an oral diabetes medication
- (e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Byetta PA

Drug Name(s)

Byetta

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

- 2. Patient had an ineffective treatment response to an oral diabetes medication
- (e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Carglumic PA

Drug Name(s)

Carglumic Acid

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA)

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of ONE of the following:

a. Acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR

b. Chronic hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR

c. Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA) AND

2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., geneticist, nephrologist, metabolic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Chenodal PA Drug Name(s) Chenodal Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require BOTH of the following: 1. Patient has a diagnosis of radiolucent stones in a well-opacifying gallbladder AND 2. The requested dose is within FDA labeled dosing for the requested indication Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Chorionic Gonadotropin PA

Drug Name(s)

Chorionic Gonadotropin

Pregnyl

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used to promote fertility AND requested agent will be used to treat erectile dysfunction AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of prepubertal cryptorchidism not due to anatomic obstruction OR

B. Patient's sex is male, with a diagnosis of hypogonadotropic hypogonadism (hypogonadism secondary to pituitary deficiency) AND BOTH of the following:

i. Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND

ii. Patient has measured luteinizing hormone (LH) AND follicle-stimulating hormone (FSH) levels that are at (low-normal) or below the testing laboratory's normal range OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Cinacalcet PA

Drug Name(s)

Cinacalcet Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

- 1. Patient has ONE of the following:
 - A. A diagnosis of hypercalcemia due to parathyroid carcinoma OR
 - B. A diagnosis of primary hyperparathyroidism (HPT) AND BOTH of the following:
 - i. Patient has a pretreatment serum calcium level that is above the testing laboratory's upper limit of normal AND
 - ii. Patient is unable to undergo parathyroidectomy OR
 - C. Another indication that is FDA approved or supported in CMS approved compendia for the requested agent not otherwise excluded from Part D [i.e., secondary hyperparathyroidism due to end-stage renal disease (ESRD) on dialysis]

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Colony Stimulating Factors PA – Fulphila

Drug Name(s)

Fulphila
Indications:
All Medically-Accepted Indications.
Off-Label Uses:
Exclusion Criteria:
Required Medical Information:
Criteria for approval require the following:
1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent
Age Restriction:
Prescriber Restrictions:
Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Granix

Drug Name(s)

Granix Indications: All Medically-Accepted Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: Prescriber Restrictions: Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious

disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Leukine

Drug Name(s)

Leukine Indications: All Medically-Accepted Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: Prescriber Restrictions: Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious

disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Nivestym

Drug Name(s)

Nivestym Indications: All Medically-Accepted Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: Prescriber Restrictions: Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious

disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Udenyca

Drug Name(s)

Udenyca

Udenyca Onbody

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Colony Stimulating Factors PA – Ziextenzo

Drug Name(s)

Ziextenzo Indications: All Medically-Accepted Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: Prescriber Restrictions: Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious

disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Corlanor PA

Drug Name(s)

Corlanor

Ivabradine

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has stable, symptomatic chronic heart failure (e.g., NYHA Class II, III, IV: ACCF/AHA Class C, D) AND

2. ONE of following:

a. ALL of the following:

i. The requested agent is for a pediatric patient, 6 months of age or over AND

ii. Patient has heart failure due to dilated cardiomyopathy (DCM) AND

iii. Patient is in sinus rhythm with an elevated heart rate OR

b. ALL of the following:

- i. The requested agent is for an adult patient AND
- ii. Patient has a baseline OR current left ventricular ejection fraction of 35% or less AND iii. Patient is in sinus rhythm with a resting heart rate of 70 beats or greater per minute prior to initiating therapy with the requested agent AND

iv. ONE of the following:

- 1. Patient is on a maximally tolerated dose of beta blocker (e.g., bisoprolol, carvedilol, metoprolol) OR
- 2. Patient has an intolerance, FDA labeled contraindications, or hypersensitivity to a beta blocker

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Cresemba PA

Drug Name(s)

Cresemba Indications: All Medically-Accepted Indications. Off-Label Uses: Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information: Criteria for initial approval require the following: 1. ONE of the following: A. Patient has a diagnosis of invasive aspergillosis OR

B. Patient has a diagnosis of invasive mucormycosis OR

C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of invasive aspergillosis and patient has continued indicators of active disease (e.g., continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR

B. Patient has a diagnosis of invasive mucormycosis and patient has continued indicators of active disease (e.g., continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR

C. BOTH of the following:

i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 6 months Other Criteria:

Crysvita PA

Drug Name(s)

Crysvita

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. ONE of the following:
 - A. BOTH of the following:

i. Patient has a diagnosis of X-linked hypophosphatemia (XLH) as confirmed by ONE of the following:

- a. Genetic testing OR
- b. Elevated levels of intact fibroblast growth factor 23 (FGF23) OR

c. Prescriber has provided information indicating the patient has a positive family history of XLH AND

- ii. ONE of the following:
 - a. Patient's epiphyseal plate has not fused OR
 - b. Patient's epiphyseal plate has fused AND the patient is experiencing symptoms of XLH (e.g., bone pain, fractures, limited mobility) OR

B. Patient has a diagnosis of tumor-induced osteomalacia (TIO) associated with phosphaturic mesenchymal tumors AND BOTH of the following:

- i. The requested agent is being used to treat FGF23 related hypophosphatemia AND
- ii. The tumor cannot be curatively surgically resected or localized AND
- 2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of X-linked hypophosphatemia (XLH) OR

B. Patient has a diagnosis of tumor-induced osteomalacia (TIO) associated with phosphaturic mesenchymal tumors AND

3. Patient has had clinical benefit with the requested agent (e.g., enhanced height velocity, improvement in lower extremity bowing and associated abnormalities, radiographic evidence of epiphyseal healing, improvement in bone pain, enhanced mobility, improvement in osteomalacia, improvement in fracture healing) AND

4. The requested dose is within FDA labeled dosing for the requested indication **Age Restriction:**

Patient is within the FDA labeled age for the requested agent for the requested indication **Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., nephrologist, endocrinologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Prior Authorization Group Description: Cystadrops PA Drug Name(s) Cystadrops Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions:** Prescriber is a specialist in the area of the patient's diagnosis (e.g., ophthalmologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Prior Authorization Group Description: Cystaran PA Drug Name(s) Cystaran Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions:** Prescriber is a specialist in the area of the patient's diagnosis (e.g., ophthalmologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Cystinosis Agents PA – Cystagon Drug Name(s)

Cystagon Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information: Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of nephropathic cystinosis AND 2. Prescriber has performed a baseline white blood cell (WBC) cystine level test AND 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of nephropathic cystinosis AND

3. Patient has had clinical benefit with the requested agent (e.g., decrease in WBC cystine levels from baseline) AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Dalfampridine PA

Drug Name(s)

Dalfampridine Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of multiple sclerosis (MS) AND

2. ONE of the following:

A. The requested agent will be used in combination with a disease modifying agent [e.g., Aubagio, Avonex, Bafiertam, Betaseron, dimethyl fumarate (e.g., Tecfidera), Extavia, fingolimod, glatiramer (e.g., Copaxone, Glatopa), Kesimpta, Mavenclad, Mayzent, Plegridy, Ponvory, Rebif, Vumerity, Zeposia] OR

B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent OR

C. Prescriber has provided information indicating that a disease modifying agent is not clinically appropriate for the patient

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of multiple sclerosis (MS) AND

3. ONE of the following:

A. The requested agent will be used in combination with a disease modifying agent [e.g., Aubagio, Avonex, Bafiertam, Betaseron, dimethyl fumarate (e.g., Tecfidera), Extavia, fingolimod, glatiramer (e.g., Copaxone, Glatopa), Kesimpta, Mavenclad, Mayzent, Plegridy, Ponvory, Rebif, Tecfidera, Vumerity, Zeposia] OR

B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent OR

C. Prescriber has provided information indicating that a disease modifying agent is not clinically appropriate for the patient AND

4. Patient has had improvements or stabilization from baseline in timed walking speed (timed 25-foot walk)

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial approval will be for 3 months, renewal approval will be for 12 months

Prior Authorization Group Description: Dayvigo PA Drug Name(s) Dayvigo Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Droxidopa PA

Drug Name(s)

Droxidopa

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of neurogenic orthostatic hypotension (nOH) AND

2. Prescriber has performed baseline blood pressure readings while the patient is sitting or supine (lying face up), AND also within three minutes of standing from a supine position AND

3. Patient has a decrease of at least 20 mmHg in systolic blood pressure or 10 mmHg diastolic blood pressure within three minutes after standing AND

4. Patient has persistent and consistent symptoms of neurogenic orthostatic hypotension (nOH) caused by ONE of the following:

A. Primary autonomic failure [Parkinson's disease (PD), multiple system atrophy, or pure autonomic failure] OR

B. Dopamine beta-hydroxylase deficiency OR

C. Non-diabetic autonomic neuropathy AND

5. Prescriber has assessed the severity of the patient's baseline symptoms of dizziness, lightheadedness, feeling faint, or feeling like the patient may black out AND

6. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of neurogenic orthostatic hypotension (nOH) AND

3. Patient has had improvements or stabilization with the requested agent as indicated by improvement in severity from baseline symptoms of ONE of the following:

A. Dizziness

B. Lightheadedness

C. Feeling faint

D. Feeling like the patient may black out AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist, neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 1 month for initial, 3 months for renewal

Dupixent PA

Drug Name(s)

Dupixent

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

- A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND ALL of the following:
 - i. ONE of the following:
 - a. Patient has tried and failed a topical steroid (e.g., triamcinolone) OR
 - b. Patient has an intolerance, hypersensitivity, or an FDA labeled
 - contraindication to a topical steroid AND
 - ii. For patients 2 years of age or over, ONE of the following:
 - a. Patient has tried and failed a topical calcineurin inhibitor (e.g., pimecrolimus, tacrolimus) OR
 - b. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to a topical calcineurin inhibitor AND
 - iii. Patient will NOT be using the requested agent in combination with another biologic agent or a JAK inhibitor for the requested indication (e.g., Adbry, Cibinqo, Opzelura, Rinvoq) OR

B. Patient has a diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma AND BOTH of the following:

i. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent AND ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication OR

C. Patient has a diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) AND the following:

i. BOTH of the following:

a. ONE of the following:

 Patient has tried and had an inadequate response to an oral systemic corticosteroid AND an intranasal corticosteroid (e.g., fluticasone) OR
 Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to an oral systemic corticosteroid AND an intranasal corticosteroid AND

Initial criteria continues: see Other Criteria Age Restriction:

For diagnosis of moderate-to-severe atopic dermatitis, patient is 6 months of age or over. For diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma, patient is 6 years of age or over. For diagnosis of CRSwNP, patient is 12 years of age or over. For diagnosis of EoE, patient is 1 year of age or over. For diagnosis of PN, patient is 18 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, dermatologist, immunologist, gastroenterologist, otolaryngologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

b. Patient will continue standard maintenance therapy (e.g., intranasal corticosteroid) in combination with the requested agent OR

- D. Patient has a diagnosis of eosinophilic esophagitis (EoE) confirmed by esophageal biopsy OR
- E. Patient has a diagnosis of prurigo nodularis (PN) AND

2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND the following:

i. Patient will NOT be using the requested agent in combination with another biologic agent or a JAK inhibitor for the requested indication (e.g., Adbry, Cibinqo, Opzelura, Rinvoq) OR

B. Patient has a diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma AND BOTH of the following:

i. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LTRA, LAMA, theophylline) in combination with the requested agent AND ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication OR

C. Patient has a diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) AND the following:

i. Patient will continue standard maintenance therapy (e.g., intranasal corticosteroid) in combination with the requested agent OR

- D. Patient has a diagnosis of eosinophilic esophagitis (EoE) OR
- E. Patient has a diagnosis of prurigo nodularis (PN) AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Emgality PA

Drug Name(s)

Emgality

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of migraine AND ALL of the following:

- i. The requested agent is being used for migraine prophylaxis AND
- ii. Patient has 4 migraine headaches or more per month AND
- iii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis OR

B. Patient has a diagnosis of episodic cluster headache AND BOTH of the following:

i. Patient has had at least 5 cluster headache attacks AND

ii. Patient has had at least two cluster periods lasting 7 days to one year and separated by pain-free remission periods of 3 months or more

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. ALL of the following:

- i. Patient has a diagnosis of migraine AND
- ii. The requested agent is being used for migraine prophylaxis AND
- iii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis OR
- B. Patient has a diagnosis of episodic cluster headache AND

3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Emsam PA

Drug Name(s)

Emsam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of major depressive disorder (MDD) OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

- C. BOTH of the following:
 - i. ONE of the following:
 - a. BOTH of the following:
 - i. Patient has a diagnosis of major depressive disorder (MDD) AND

ii. ONE of the following:

1. Patient has tried and had an inadequate response to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR

2. Patient has an intolerance or hypersensitivity to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR

3. Patient has an FDA labeled contraindication to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of major depressive disorder (MDD) OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days

OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. BOTH of the following:

i. Patient has had clinical benefit with the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Endari PA

Drug Name(s)

Endari

L-glutamine

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of sickle cell disease AND
- 2. Patient is using the requested agent to reduce the acute complications of sickle cell disease AND
- 3. ONE of the following:

A. Patient has tried and had an inadequate response to maximally tolerated dose of hydroxyurea OR

- B. Patient has an intolerance or hypersensitivity to hydroxyurea OR
- C. Patient has an FDA labeled contraindication to hydroxyurea AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration: Approval will be for 12 months

Epclusa PA

Drug Name(s)

Epclusa

Sofosbuvir/Velpatasvir

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of hepatitis C confirmed by serological markers OR

B. Patient is a hepatitis C virus (HCV) - uninfected solid organ transplant recipient AND BOTH of the following:

i. Patient received an HCV - viremic donor organ AND

ii. The requested agent is being used for prophylaxis AND

2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND

3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND

4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND

5. ONE of the following:

A. The requested agent is the preferred agent: Epclusa OR

B. The requested agent is the non-preferred agent: sofosbuvir/velpatasvir AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR

iv. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration:**

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Epidiolex PA

Drug Name(s)

Epidiolex

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of seizures associated with ONE of the following:

A. Lennox-Gastaut syndrome OR

- B. Dravet syndrome OR
- C. Tuberous sclerosis complex AND

2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Erythropoietin Stimulating Agents PA – Aranesp

Drug Name(s)

Aranesp Albumin Free

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. The intent of chemotherapy is non-curative OR

B. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

C. Anemia due to myelodysplastic syndrome AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR

D. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restriction:

Prescriber Restrictions: Coverage Duration: 6 months for chemotherapy, 12 months for other indications Other Criteria:

Erythropoietin Stimulating Agents PA - Epogen/Procrit

Drug Name(s)

Procrit

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR

B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. The intent of chemotherapy is non-curative OR

C. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

D. Anemia due to myelodysplastic syndrome AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR

E. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other

Other Criteria:

F. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Erythropoietin Stimulating Agents PA – Retacrit

Drug Name(s)

Retacrit

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR

B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. The intent of chemotherapy is non-curative OR

C. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks)

AND

ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

D. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR

E. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review. Age Restriction:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other **Other Criteria:**

Prior Authorization Group Description: Eysuvis PA Drug Name(s) Eysuvis Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require ALL of the following: 1. Patient has a diagnosis of dry eye disease AND 2. The requested agent will be used for short-term (up to two weeks) treatment AND 3. The requested dose is within FDA labeled dosing for the requested indication Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 1 month **Other Criteria:**

Fasenra PA

Drug Name(s)

Fasenra

Fasenra Pen Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of severe asthma with an eosinophilic phenotype AND

2. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA,

LTRA, LAMA, theophylline) in combination with the requested agent AND

3. Patient will NOT be using the requested agent in combination with Xolair, Dupixent, or with another injectable interleukin 5 (IL-5) inhibitor (e.g., Cingair, Nucala) for the requested indication AND

4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of severe asthma with an eosinophilic phenotype AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA,

LTRA, LAMA, theophylline) in combination with the requested agent AND

5. Patient will NOT be using the requested agent in combination with Xolair, Dupixent, or with another injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Nucala) for the requested indication AND

6. The requested dose is within the FDA labeled dosing for the requested indication

Age Restriction:

Patient is 6 years of age or over

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist,

pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration**:

Approval will be for 12 months **Other Criteria**:

Fentanyl Oral PA - Fentanyl lozenge

Drug Name(s)

Fentanyl Citrate Oral Transmucosal

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. ONE of the following:

a. Patient has a documented diagnosis (i.e., medical records) of chronic cancer pain due to an active malignancy AND the following:

i. There is evidence of a claim that the patient is currently being treated with a longacting opioid with the requested agent within the past 90 days OR

b. Patient has a diagnosis that is supported in CMS approved compendia for the requested agent AND

2. Patient will NOT be using the requested agent in combination with any other oral or nasal fentanyl agent

Age Restriction:

Patient is 16 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Fintepla PA

Drug Name(s)

Fintepla

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of seizures associated with Dravet syndrome (DS) or Lennox-Gastaut syndrome (LGS) AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. An echocardiogram assessment will be obtained before and during treatment with the requested agent, to evaluate for valvular heart disease and pulmonary arterial hypertension AND

ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND

iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Focalin PA Drug Name(s) Dexmethylphenidate Hydrochloride Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Gammagard/Gammaked/Gamunex-C PA

Drug Name(s)

Gammagard Liquid

Gammagard S/D Iga Less Than 1Mcg/MI

Gamunex-C

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ONE of the following:

1. Patient has ONE of the following diagnoses:

A. Primary immunodeficiency [e.g., congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, Xlinked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency] OR

B. B-cell chronic lymphocytic leukemia OR multiple myeloma AND ONE of the following:

i. Patient has a history of infections OR

ii. Patient has evidence of specific antibody deficiency OR

iii. Patient has hypogammaglobulinemia OR

C. Idiopathic thrombocytopenia purpura AND ONE of the following:

i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g.,

methylprednisolone), or immunosuppressants (e.g., azathioprine)] OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

D. Dermatomyositis AND ONE of the following:

i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g.,

methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

E. Polymyositis AND ONE of the following:

i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g.,

methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

F. Severe rheumatoid arthritis AND ONE of the following:

i. Patient has failed ONE conventional therapy [e.g., tumor necrosis factor antagonists (e.g., Humira), DMARDS (e.g., methotrexate)] OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 6 months for indications in Other Criteria, 12 months for all others **Other Criteria**:

G. Myasthenia gravis (MG) AND ONE of the following:

i. Patient is in acute myasthenic crisis OR

ii. Patient has severe refractory MG (e.g., major functional disability/weakness) AND ONE of the following:

a) Patient has failed ONE immunomodulator therapy (i.e., corticosteroid, pyridostigmine, or azathioprine) OR

b) Patient has an intolerance, FDA labeled contraindication, or hypersensitivity

to ONE immunomodulator therapy OR

H. Multiple sclerosis (MS) AND BOTH of the following:

i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND

ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication to TWO MS agents (e.g., Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR

I. Acquired von Willebrand hemophilia AND ONE of the following:

i. Patient has failed ONE conventional therapy (e.g., desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, or recombinant factor VIIa) OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

J. Refractory pemphigus vulgaris AND ONE of the following:

i. Patient has failed ONE conventional immunosuppressive therapy (e.g., azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR

ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR

2. ONE of the following:

A. Patient has another FDA labeled indication for the requested agent OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 6 months approval duration: Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcus, Toxic epidermal necrolysis and Stevens-Johnson syndrome

Drug is also subject to Part B versus Part D review.

Gattex PA

Drug Name(s)

Gattex

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of short bowel syndrome (SBS) AND
- 2. Patient is dependent on parenteral nutrition OR intravenous (PN/IV) fluids AND
- 3. ONE of the following:
 - A. Patient is aged 1 year to 17 years AND BOTH of the following:
 - i. A fecal occult blood test has been performed within 6 months prior to initiating treatment with the requested agent AND
 - ii. ONE of the following:
 - a. There was no unexplained blood in the stool OR
 - b. There was unexplained blood in the stool AND a colonoscopy or a sigmoidoscopy was performed OR
 - B. Patient is 18 years of age or over AND BOTH of the following:
 - i. Patient has had a colonoscopy within 6 months prior to initiating treatment with the requested agent AND
 - ii. If polyps were present at this colonoscopy, the polyps were removed AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of short bowel syndrome (SBS) AND

3. Patient has had a reduction from baseline in parenteral nutrition OR intravenous (PN/IV) fluids AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal **Other Criteria:**

Gaucher Enzyme Replacement PA – Cerezyme

Drug Name(s)

Cerezyme

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:

A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR

B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND

2. Prescriber has drawn baseline measurements of hemoglobin, platelet count, liver volume, and spleen volume AND

3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:

A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR

- B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
- C. Hepatomegaly OR
- D. Splenomegaly OR
- E. Growth failure (i.e., growth velocity below the standard mean for age) OR
- F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND

- 3. Patient has had improvement and/or stabilization from baseline in at least ONE of the following:
 - A. Hemoglobin (Hb) level OR
 - B. Platelet count OR
 - C. Liver volume OR
 - D. Spleen volume OR
 - E. Growth velocity OR
 - F. Bone pain or disease

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist,

hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Gaucher Enzyme Replacement PA – Elelyso

Drug Name(s)

Elelyso

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:

A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR

B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND

2. Prescriber has drawn baseline measurements of hemoglobin, platelet count, liver volume, and spleen volume AND

3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:

A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR

- B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
- C. Hepatomegaly OR
- D. Splenomegaly OR
- E. Growth failure (i.e., growth velocity below the standard mean for age) OR
- F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND

- 3. Patient has had improvement and/or stabilization from baseline in at least ONE of the following:
 - A. Hemoglobin (Hb) level OR
 - B. Platelet count OR
 - C. Liver volume OR
 - D. Spleen volume OR
 - E. Growth velocity OR
 - F. Bone pain or disease

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist,

hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Gaucher Enzyme Replacement PA – Vpriv

Drug Name(s)

Vpriv

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:

A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR

B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND

2. Prescriber has drawn baseline measurements of hemoglobin, platelet count, liver volume, and spleen volume AND

3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:

A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR

- B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
- C. Hepatomegaly OR
- D. Splenomegaly OR
- E. Growth failure (i.e., growth velocity below the standard mean for age) OR
- F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND

- 3. Patient has had improvement and/or stabilization from baseline in at least ONE of the following:
 - A. Hemoglobin (Hb) level OR
 - B. Platelet count OR
 - C. Liver volume OR
 - D. Spleen volume OR
 - E. Growth velocity OR
 - F. Bone pain or disease

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist,

hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Growth Hormone PA – Omnitrope

Drug Name(s)

Omnitrope

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

For Children – Criteria for initial approval require the following:

1. ONE of the following:

- a. Patient has a diagnosis of Turner Syndrome OR
- b. Patient has a diagnosis of Prader-Willi Syndrome OR
- c. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:

i. Deficiencies in 3 or more pituitary axes AND

ii. Measured serum IGF-1 (insulin-like growth factor-1) levels are below the age and sexappropriate reference range when off GH therapy OR

d. Patient has a diagnosis of growth hormone deficiency (GHD) or short stature AND BOTH of the following:

i. Patient has ONE of the following:

a) Height more than 2 standard deviations (SD) below the mean for age and sex OR

b) Height more than 1.5 SD below the midparental height OR

c) A decrease in height SD of more than 0.5 over one year in children at least 2 years of age OR

d) Height velocity more than 2 SD below the mean over one year or more than 1.5 SD sustained over two years AND

ii. Failure of at least 2 growth hormone (GH) stimulation tests (e.g., peak GH value of less than 10 mcg/L after stimulation, or otherwise considered abnormal as determined by testing lab) OR

e. Patient has a diagnosis of small for gestational age (SGA) AND ALL of the following:

i. Patient is at least 2 years of age AND

ii. Documented birth weight and/or length that is 2 or more SD below the mean for gestational age AND

iii. At 24 months of age, the patient fails to manifest catch-up growth evidenced by a height that remains 2 or more SD below the mean for age and sex

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

For Children – Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the preferred agent through the plan's Prior Authorization criteria AND

2. Patient has been diagnosed with ONE of the following:

- a. Growth Hormone Deficiency, Short Stature OR
- b. Panhypopituitarism OR
- c. Prader-Willi Syndrome OR
- d. Small for Gestational Age (SGA) OR
- e. Turner Syndrome AND

3. ALL of the following:

- a. Patient does NOT have closed epiphyses AND
- b. Patient is being monitored for adverse effects of therapy with the requested agent AND
- c. Patient's height has increased or height velocity has improved since initiation or last approval
- of the requested agent

For Adults – Criteria for initial approval require the following:

1. Patient has been diagnosed with ONE of the following:

a. Childhood growth hormone deficiency (GHD) with genetic or organic origin AND ONE of the following:

i. Low IGF-1 (insulin-like growth factor-1) level without GH replacement therapy OR ii. Failure of at least one growth hormone (GH) stimulation test as an adult (e.g., peak GH value of 5 mcg/L or lower after stimulation, or otherwise considered abnormal as determined by testing lab) OR

b. Acquired adult GHD secondary to structural lesions or trauma AND ONE of the following:

i. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:

a) Deficiencies in 3 or more pituitary axes AND

b) Low IGF-1 level without GH replacement therapy OR

ii. Patient has failed at least one growth hormone (GH) stimulation test as an adult OR c. Idiopathic GHD (adult or childhood onset) AND the patient has failed at least two growth hormone (GH) stimulation tests as an adult

For Adults – Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the preferred agent through the plan's Prior Authorization criteria AND

2. Patient has been diagnosed with ONE of the following:

- a. Childhood growth hormone deficiency (GHD) with genetic or organic origin OR
- b. Acquired adult GHD secondary to structural lesions or trauma OR
- c. Idiopathic GHD (adult or childhood onset) AND
- 3. Patient is being monitored for adverse effects of therapy with the requested agent AND
- 4. Patient's IGF-1 level has been evaluated to confirm the appropriateness of the current dose AND

5. Patient has had clinical benefit with the requested agent (i.e., body composition, hip-to-waist ratio, cardiovascular health, bone mineral density, serum cholesterol, physical strength, or quality of life)

HAE PA – Cinryze

Drug Name(s)

Cinryze Indications: All FDA-Approved Indications, Some Medically-Accepted Indications. Off-Label Uses: Acute HAE attacks Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:

a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR

b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR

c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:

i. BOTH of the following:

1. Family history of angioedema AND

2. ALL other causes of angioedema have been ruled out OR

ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation,plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase6 gene mutation, or myoferlin gene mutation that is associated with the disease AND

2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND

3. ONE of the following:

a. The requested agent will be used to treat acute HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks OR

b. The requested agent will be used for prophylaxis against HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria

AND

2. Patient has a diagnosis of hereditary angioedema (HAE) AND ONE of the following:

a. The requested agent will be used to treat acute HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks OR

b. The requested agent will be used for prophylaxis against HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks AND

3. Patient has had a decrease in the frequency or severity of acute attacks or has had stabilization of disease from use of the requested agent

HAE PA – Haegarda

Drug Name(s)

Haegarda

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:

a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR

b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR

c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:

i. BOTH of the following:

1. Family history of angioedema AND

2. ALL other causes of angioedema have been ruled out OR

ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND

2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND

3. The requested agent will be used for prophylaxis against HAE attacks AND

4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of hereditary angioedema (HAE) AND

3. The requested agent is being used for prophylaxis against HAE attacks AND

4. Patient has had a decrease in the frequency or severity of acute attacks or has had stabilization of disease from use of the requested agent AND

5. Patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

HAE PA – Icatibant

Drug Name(s)

Icatibant Acetate

Sajazir

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:

a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR

b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR

c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:

i. BOTH of the following:

1. Family history of angioedema AND

2. ALL other causes of angioedema have been ruled out OR

ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND

2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND

3. The requested agent will be used to treat acute HAE attacks AND

4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of hereditary angioedema (HAE) AND

3. The requested agent will be used to treat acute HAE attacks AND

4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks AND

5. Patient has had a decrease in the frequency or severity of acute attacks or stabilization of disease from use of the requested agent

Harvoni PA

Drug Name(s)

Harvoni

Ledipasvir/Sofosbuvir

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND

2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND

The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
 The requested dose is within FDA labeled dosing or supported in AASLD/IDSA dosing for the

- requested indication AND
- 5. ONE of the following:
 - A. The requested agent is the preferred agent: Harvoni OR

B. The requested agent is the non-preferred agent: ledipasvir/sofosbuvir AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR

iv. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria**:

High Risk Medication PA - All Starts

Drug Name(s)

Benztropine Mesylate

Clemastine Fumarate

Cyproheptadine Hydrochloride

Dicyclomine Hydrochloride

Diphenoxylate Hydrochloride/Atropine Sulfate

Hydroxyzine Hydrochloride

Hydroxyzine Pamoate

Methscopolamine Bromide

Promethazine Hcl Plain

Promethazine Hydrochloride

Promethegan

Scopolamine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients less than 65 years of age.

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high-risk medication AND

2. Prescriber has indicated that the benefits of the requested high-risk medication outweigh the risks for the patient AND

3. Prescriber has indicated that the risks and potential side effects of the requested high-risk medication have been discussed with the patient

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Ilaris PA

Drug Name(s)

llaris

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute gouty arthritis

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of:

1. ONE of the following:

A. Patient (pt) has been diagnosed with Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) OR

B. Pt has been diagnosed with Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) OR

C. Pt has been diagnosed with Familial Mediterranean Fever (FMF) AND ONE of the following:

i. Pt has tried and had an inadequate response to colchicine OR

ii. Pt has an intolerance or hypersensitivity to colchicine OR

iii. Pt has an FDA labeled contraindication to colchicine OR

D. Pt has been diagnosed with Cryopyrin-Associated Periodic Syndrome (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR

E. Pt has been diagnosed with active Systemic Juvenile Idiopathic Arthritis (SJIA) AND BOTH of the following:

i. Pt has documented active systemic features (e.g., ongoing fever, evanescent erythematous rash, generalized lymphadenopathy, 1 or more joints with active arthritis, hepatomegaly, splenomegaly, serositis) AND

ii. ONE of the following:

a. Pt has tried and had an inadequate response to at least ONE prerequisite agent (e.g., glucocorticosteroids, prescription oral NSAIDs, methotrexate, leflunomide) OR

b. Pt has an intolerance or hypersensitivity to at least ONE prerequisite agent OR

c. Pt has an FDA labeled contraindication to at least ONE prerequisite agent OR F. Pt has a diagnosis of adult onset Still's disease OR

G. Pt has been diagnosed with acute gouty arthritis AND ONE of the following:

i. Pt has tried and had an inadequate response to at least TWO conventional first-line agents (e.g., prescription oral NSAIDs, colchicine, systemic corticosteroids) OR

ii. Pt has an intolerance or hypersensitivity to at least TWO conventional first-line agents OR

iii. Pt has an FDA labeled contraindication to at least TWO conventional first-line agents AND

2. Pt will NOT be using the requested agent in combination with another biologic agent

Age Restriction:

For diagnosis of CAPS including FCAS or MWS, patient is 4 years of age or over

For diagnosis of SJIA, patient is 2 years of age or over **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Imiquimod PA

Drug Name(s)

Imiquimod

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. Patient has ONE of the following diagnoses:

- A. Actinic keratosis OR
- B. Superficial basal cell carcinoma OR
- C. External genital and/or perianal warts/condyloma acuminata OR
- D. Squamous cell carcinoma OR
- E. Basal cell carcinoma OR
- F. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

4 months for Actinic keratosis, other diagnoses - see Other Criteria

Other Criteria:

2 months for Superficial basal cell carcinoma, Squamous cell carcinoma, or Basal cell carcinoma

4 months for External genital and/or perianal warts/condyloma acuminata

12 months for All other diagnoses

Inbrija PA

Drug Name(s)

Inbrija

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. The requested agent will be used for intermittent treatment of OFF episodes in patients with Parkinson's disease AND

2. Patient is receiving concurrent therapy with carbidopa/levodopa AND

3. Patient will NOT be using a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine,

tranylcypromine) in combination with, or within 2 weeks of, the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Injectable Oncology PA Drug Name(s) Abraxane Adcetris Aliqopa Alymsys Avastin Beleodaq Besponsa Blenrep Blincyto Bortezomib Cyramza Danyelza Darzalex Darzalex Faspro Doxorubicin Hydrochloride Liposomal Empliciti Enhertu Erbitux Eribulin Folotyn Fulvestrant Gazyva Halaven Herceptin Herceptin Hylecta Herzuma Jevtana Kadcyla Kanjinti **Kyprolis** Lumoxiti Margenza Monjuvi Mvasi

Mylotarg

Nelarabine

Ogivri

Onivyde

Ontruzant

Padcev

Pemetrexed

Perjeta

Phesgo

Polivy

Portrazza

Poteligeo

Romidepsin

Rybrevant

Sarclisa

Synribo

Trazimera

Trodelvy

Unituxin

Vectibix

Vegzelma

Vyxeos

Yondelis

Zaltrap

Zepzelca

Zirabev

Zynlonta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia

for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported by CMS approved compendia as first-line therapy for the requested indication OR
b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested indication OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled contraindications to the requested agent AND

iv. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

May also be subject to Part B versus Part D review.

Age Restriction: Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Iron Chelating Agents PA – Exjade

Drug Name(s)

Deferasirox (Exjade)

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:

i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR

ii. A serum ferritin greater than 300 mcg/L OR

iii. MRI confirmation of iron deposition OR

B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND

2. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR

- B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
- 3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Iron Chelating Agents PA – Jadenu

Drug Name(s)

Deferasirox (Jadenu)

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:

i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR

ii. A serum ferritin greater than 300 mcg/L OR

iii. MRI confirmation of iron deposition OR

B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND

2. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR

- B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
- 3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Ivermectin Cream PA Drug Name(s) Ivermectin Cream Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Ivermectin Tablet PA

Drug Name(s)

Ivermectin Tablet

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 4 months

Kalydeco PA

Drug Name(s)

Kalydeco Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of cystic fibrosis AND

2. ONE of the following:

A. Patient has ONE of the CFTR gene mutations or a mutation in the CFTR gene that is responsive based on in vitro data, as indicated in the FDA label, confirmed by genetic testing ORB. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as

indicated in the FDA label, confirmed by genetic testing AND

3. Patient is NOT homozygous for the F508del mutation AND

4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of cystic fibrosis AND

Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
 Patient will NOT be using the requested agent in combination with another CFTR modulator agent for

the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Kerendia PA Drug Name(s) Kerendia Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Korlym PA

Drug Name(s)

Korlym

Mifepristone

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

- Criteria for initial approval require ALL of the following:
- 1. Patient has a diagnosis of Cushing's syndrome AND
- 2. ONE of the following:
 - A. Patient has type 2 diabetes mellitus OR

B. Patient has glucose intolerance as defined by a 2-hour glucose tolerance test plasma glucose value of 140-199 mg/dL AND

- 3. ONE of the following:
 - A. Patient has failed surgical resection OR
 - B. Patient is NOT a candidate for surgical resection

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of Cushing's syndrome AND

3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Leuprolide PA

Drug Name(s)

Eligard

Leuprolide Acetate

Lupron Depot (1-Month)

Lupron Depot (3-Month)

Lupron Depot (4-Month)

Lupron Depot (6-Month)

Lupron Depot-Ped (1-Month)

Lupron Depot-Ped (3-Month)

Lupron Depot-Ped (6-Month)

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

- C. BOTH of the following:
 - i. Patient is NOT currently being treated with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND

3. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Lidocaine Topical PA - Lidocaine Gel/Jelly

Drug Name(s)

Glydo

Lidocaine Hcl Gel

Lidocaine Hcl Jelly

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:

- A. Surface anesthesia and lubrication for urethral procedure OR
- B. Topical treatment for pain of urethritis OR
- C. Surface anesthesia and lubrication for endotracheal intubation (oral and nasal) OR

D. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine Ointment

Drug Name(s)

Lidocaine Ointment

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:

- A. Anesthesia of accessible mucous membranes of the oropharynx OR
- B. Anesthetic lubricant for intubation OR

C. Temporary relief of pain associated with minor burns, including sunburn, abrasions of the skin, and insect bites OR

D. Another indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

i. Patient has tried and had an inadequate response to a conventional therapy [e.g., gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR

ii. Patient has an intolerance or hypersensitivity to a conventional therapy OR

iii. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Lidocaine Topical PA - Lidocaine Patch

Drug Name(s)

Lidocan

Lidocan III

Lidocaine Patch

Tridacaine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:

- A. Pain associated with postherpetic neuralgia (PHN) OR
- B. Pain associated with diabetic neuropathy OR
- C. Neuropathic pain associated with cancer, or cancer treatment OR

D. Another diagnosis that is supported in CMS approved compendia for the requested agent AND

- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to a conventional therapy [e.g.,

gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR

B. Patient has an intolerance or hypersensitivity to a conventional therapy OR

C. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine Solution

Drug Name(s)

Lidocaine Solution

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:

A. Topical anesthesia of accessible mucous membranes of the oral and nasal cavities OR

B. Topical anesthesia of accessible mucous membranes of proximal portions of the digestive tract OR

C. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA - Lidocaine/prilocaine Cream

Drug Name(s)

Lidocaine/Prilocaine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:

- A. Local analgesia on normal intact skin OR
- B. Topical anesthetic for dermal procedures OR
- C. Adjunctive anesthesia prior to local anesthetic infiltration in adult male genital skin OR
- D. Anesthesia for minor procedures on female external genitalia OR
- E. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Lidocaine Topical PA – ZTlido

Drug Name(s)

Ztlido

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

- A. Pain associated with postherpetic neuralgia (PHN) OR
- B. Neuropathic pain associated with cancer, or cancer treatment OR
- C. Another diagnosis that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. Patient has tried and had an inadequate response to generic lidocaine 5% patch OR
- B. Patient has an intolerance or hypersensitivity to generic lidocaine 5% patch OR
- C. Patient has an FDA labeled contraindication to generic lidocaine 5% patch AND
- 3. ONE of the following:

A. Patient has tried and had an inadequate response to a conventional therapy [e.g., gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR

- B. Patient has an intolerance or hypersensitivity to a conventional therapy OR
- C. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Linezolid PA

Drug Name(s)

Linezolid

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND ONE of the following:

a. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR

b. Patient has a documented infection due to vancomycin-resistant Enterococcus faecium OR

c. Patient has a diagnosis of pneumonia caused by Staphylococcus aureus or Streptococcus pneumoniae AND ONE of the following:

i. Patient has a documented infection that is resistant to TWO of the following: betalactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole, OR that is resistant to vancomycin OR

ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

- iv. Patient has an intolerance or hypersensitivity to vancomycin OR
- v. Patient has an FDA labeled contraindication to vancomycin OR

d. Patient has a documented skin and skin structure infection, including diabetic foot infections, caused by Staphylococcus aureus, Streptococcus pyogenes, or Streptococcus agalactiae AND ONE of the following:

i. Patient has a documented infection that is resistant to TWO of the following: betalactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole, OR that is resistant to vancomycin at the site of infection OR

ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 3 months

Other Criteria:

iv. Patient has an intolerance or hypersensitivity to vancomycin OR

v. Patient has an FDA labeled contraindication to vancomycin AND

2. Patient will NOT be using the requested agent in combination with Sivextro (tedizolid) for the same infection AND

3. The requested dose is within FDA labeled dosing for the requested indication

Lumryz PA

Drug Name(s)

Lumryz

Lumryz Pak

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of narcolepsy with cataplexy OR

B. Patient has a diagnosis of narcolepsy with excessive daytime sleepiness AND BOTH of the following:

i. ONE of the following:

a. Patient has tried and had an inadequate response to modafinil or armodafinil OR

b. Patient has an intolerance or hypersensitivity to modafinil or armodafinil OR

c. Patient has an FDA labeled contraindication to modafinil or armodafinil AND

ii. ONE of the following:

a. Patient has tried and had an inadequate response to ONE standard stimulant agent (e.g., methylphenidate) OR

b. Patient has an intolerance or hypersensitivity to ONE standard stimulant agent (e.g., methylphenidate) OR

c. Patient has an FDA labeled contraindication to ONE standard stimulant agent (e.g., methylphenidate) OR

C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Patient is 18 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Memantine ER PA

Drug Name(s)

Memantine Hydrochloride Er

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients greater than or equal to 30 years of age

Criteria for approval require the following:

1. Patient is younger than 30 years of age AND ONE of the following:

- A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Memantine PA

Drug Name(s)

Memantine Hcl Titration Pak

Memantine Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

PA does NOT apply to patients greater than or equal to 30 years of age

Criteria for approval require the following:

1. Patient is younger than 30 years of age AND ONE of the following:

A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Methylin PA Drug Name(s) Methylphenidate Hydrochloride (Methylin) Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Methylphenidate ER Tablet PA Drug Name(s)

Methylphenidate Hydrochloride Er Tablet Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Migranal PA

Drug Name(s)

Dihydroergotamine Mesylate Spray

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. The requested agent will be used for the treatment of acute migraine with or without aura AND

2. ONE of the following:

A. Patient has tried and had an inadequate response to TWO triptan agents with differing active ingredients (e.g., sumatriptan, rizatriptan) OR

B. Patient has an intolerance or hypersensitivity to TWO triptan agents with differing active ingredients OR

C. Patient has an FDA labeled contraindication to TWO triptan agents with differing active ingredients AND

3. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, acute CGRP)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. The requested agent will be used for the treatment of acute migraine with or without aura AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, acute CGRP)

Age Restriction:

Prescriber Restrictions: Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Modafinil PA Drug Name(s) Modafinil Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require BOTH of the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another target agent (i.e., armodafinil) Age Restriction: Patient is 17 years of age or over **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Mounjaro PA

Drug Name(s)

Mounjaro

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has a diagnosis of type 2 diabetes mellitus AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

2. Patient had an ineffective treatment response to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

MS PA – Avonex

Drug Name(s)

Avonex

Avonex Pen Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information: Criteria for initial approval require BOTH of the following: 1. Patient has an FDA labeled indication for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication Criteria for renewal approval require ALL of the following: 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND 2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent

(DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Betaseron Drug Name(s) Betaseron Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information: Criteria for initial approval require BOTH of the following: 1. Patient has an FDA labeled indication for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

MS PA - Dimethyl Fumarate

Drug Name(s)

Dimethyl Fumarate

Dimethyl Fumarate Starterpack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND

- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent
- (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent

(DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA - Fingolimod Drug Name(s) Fingolimod Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for initial approval require ALL of the following: 1. Patient has an FDA labeled indication for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication AND 3. Prescriber has performed an electrocardiogram within 6 months prior to initiating treatment Criteria for renewal approval require ALL of the following: 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND 2. Patient has an FDA labeled indication for the requested agent AND 3. Patient has had clinical benefit with the requested agent AND 4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication Age Restriction: **Prescriber Restrictions:**

Coverage Duration:

Approval will be for 12 months Other Criteria: **Prior Authorization Group Description:** MS PA – Glatiramer Drug Name(s) Copaxone Glatiramer Acetate Glatopa Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for initial approval require BOTH of the following: 1. Patient has an FDA labeled indication for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication Criteria for renewal approval require ALL of the following: 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND 2. Patient has an FDA labeled indication for the requested agent AND 3. Patient has had clinical benefit with the requested agent AND 4. Patient will NOT be using the requested agent in combination with another disease modifying agent

4. Patient will NOT be using the requested agent in combination with another disease modifying ag (DMA) or a biologic immunomodulator for the requested indication

Age Restriction: Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

MS PA – Kesimpta Drug Name(s) Kesimpta Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: FDA labeled contraindications to the requested agent Required Medical Information: Criteria for initial approval require BOTH of the following: 1. Patient has an FDA labeled indication for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

MS PA – Mayzent

Drug Name(s)

Mayzent

Mayzent Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND

2. Patient will NOT be using the requested agent in combination with another disease modifying agent

(DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent

(DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration: Approval will be for 12 months

MS PA – Plegridy

Drug Name(s)

Plegridy

Plegridy Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND

- 2. Patient will NOT be using the requested agent in combination with another disease modifying agent
- (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent

(DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

MS PA – Tysabri

Drug Name(s)

Tysabri Indications: All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. Patient has ONE of the following diagnoses:

i. Relapsing form of Multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, or active secondary progressive disease AND ONE of the following:

a. Prescriber states the patient has highly active disease OR

b. ONE of the following:

 Patient has tried and had an inadequate response to TWO preferred agents (Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR
 Patient has an intolerance or hypersensitivity to TWO preferred agents (Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR
 Patient has an FDA labeled contraindication to TWO preferred agents (Avonex, Betaseron, Copaxone, dimethyl fumarate, fingolimod, glatiramer, Glatopa, Mayzent, Plegridy, Vumerity) OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months for MS, for CD 16 weeks for initial and 12 months renewal **Other Criteria:**

ii. Moderately to severely active Crohn's disease (CD) AND BOTH of the following:

a. ONE of the following:

1. Patient has tried and had an inadequate response to ONE conventional CD therapy (e.g., 6-mercaptopurine, azathioprine, corticosteroids, methotrexate) OR

2. Patient has an intolerance or hypersensitivity to ONE conventional CD therapy (e.g., 6-mercaptopurine, azathioprine, corticosteroids, methotrexate) OR

3. Patient has an FDA labeled contraindication to ONE conventional CD therapy (e.g., 6-mercaptopurine, azathioprine, corticosteroids, methotrexate) AND

b. ONE of the following:

 Patient has tried and had an inadequate response to ONE preferred biologic agent (Humira, Skyrizi, or Stelara) for the treatment of CD OR
 Patient has an intolerance or hypersensitivity to ONE preferred biologic agent (Humira, Skyrizi, or Stelara) OR

3. Patient has an FDA labeled contraindication to ONE preferred biologic agent (Humira, Skyrizi, or Stelara) AND

3. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent

(DMA) or a biologic immunomodulator for the requested indication

MS PA – Vumerity **Drug Name(s)** Vumerity **Indications:** All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for initial approval require BOTH of the following: 1. Patient has an FDA labeled indication for the requested agent AND 2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Myalept PA

Drug Name(s)

Myalept Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has leptin deficiency associated with a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND

2. Prescriber has provided the patient's baseline levels for HbA1C, triglycerides, and fasting insulin, measured prior to beginning therapy with the requested agent AND

 Patient also has at least ONE of the complications related to lipodystrophy: diabetes mellitus, hypertriglyceridemia (200 mg/dL or higher), and/or high fasting insulin (30μU/mL or higher) AND
 Patient has tried and had an inadequate response to maximum tolerable dosing of a conventional agent for the additional diagnosis AND

5. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has leptin deficiency associated with a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND

3. Patient has had improvement or stabilization with the requested agent as indicated by change from baseline level of at least ONE of the following:

- A. HbA1C
- B. Triglycerides
- C. Fasting insulin AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Conventional agent examples include:

Hypertriglyceridemia: statins, fenofibrates, Omega-3-Acid Ethyl Esters (generic Lovaza)

Diabetes/high fasting insulin: insulin, sulfonylurea/sulfonylurea combination, metformin/metformin combination

Natpara PA

Drug Name(s)

Natpara Indications: All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Increased baseline risk for osteosarcoma (e.g., Paget's disease of bone, unexplained elevations of alkaline phosphatase, hereditary disorders predisposing to osteosarcoma, history of external beam or implant radiation therapy involving the skeleton, pediatric and young adult patients with open epiphyses)

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hypocalcemia associated with hypoparathyroidism AND

2. Patient does NOT have a baseline vitamin D level below the testing laboratory's lower limit of normal AND

3. Patient's baseline serum calcium level (albumin-corrected) is above 7.5 mg/dL AND

4. Patient will NOT be using the requested agent in combination with alendronate for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of hypocalcemia associated with hypoparathyroidism AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with alendronate for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, nephrologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Nuedexta PA

Drug Name(s)

Nuedexta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of pseudobulbar affect OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) [e.g., Marplan (isocarboxazid), Nardil (phenelzine), and Parnate (tranylcypromine)]

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Prior Authorization Group Description: Nuplazid PA Drug Name(s) Nuplazid Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria: Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Nurtec PA

Drug Name(s)

Nurtec

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

- 1. Patient has a diagnosis of migraine AND
- 2. ONE of the following:

A. The requested agent is being used for the treatment of acute migraine with or without aura AND BOTH of the following:

i. ONE of the following:

a. Patient has tried and had an inadequate response to a triptan (e.g.,

sumatriptan, rizatriptan) agent OR

- b. Patient has an intolerance, or hypersensitivity to a triptan OR
- c. Patient has an FDA labeled contraindication to a triptan AND

ii. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP) OR

B. The requested agent is being used for migraine prophylaxis AND BOTH of the following:

i. Patient has 4 migraine headaches or more per month AND

- ii. Patient will NOT be using the requested agent in combination with another calcitonin
- gene-related peptide (CGRP) agent for migraine prophylaxis

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 Months

Other Criteria:

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has a diagnosis of migraine AND
- 3. ONE of the following:

A. The requested agent is being used for the treatment of acute migraine with or without aura AND BOTH of the following:

i. Patient has had clinical benefit with the requested agent AND

ii. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP) OR

B. The requested agent is being used for migraine prophylaxis AND BOTH of the following:

i. Patient has had clinical benefit with the requested agent AND

ii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Ocaliva PA

Drug Name(s)

Ocaliva

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of primary biliary cholangitis (PBC) confirmed by at least TWO of the following:
 - A. There is biochemical evidence of cholestasis with an alkaline phosphatase (ALP) elevation
 - B. Presence of antimitochondrial antibody (AMA): a titer greater than 1:80 OR a level that is above the testing laboratory's upper limit of the normal range

C. If the AMA is negative or present only in low titer (less than or equal to 1:80), presence of other PBC-specific autoantibodies, including sp100 or gp210

D. Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND

- 2. ONE of the following:
 - A. Patient does NOT have cirrhosis OR
 - B. Patient has compensated cirrhosis with NO evidence of portal hypertension AND
- 3. Prescriber has measured the patient's alkaline phosphatase (ALP) level AND total bilirubin level AND
- 4. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has tried and had an inadequate response to ursodiol AND
 - ii. The requested agent will be used in combination with ursodiol OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has a diagnosis of primary biliary cholangitis (PBC) AND
- 3. ONE of the following:
 - A. Patient does NOT have cirrhosis OR
 - B. Patient has compensated cirrhosis with NO evidence of portal hypertension AND
- 4. ONE of the following:
 - A. The requested agent will be used in combination with ursodiol OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol AND

5. Patient has had improvements or stabilization with the requested agent as indicated by BOTH of the following:

- A. Decrease in alkaline phosphatase (ALP) level from baseline AND
- B. Total bilirubin is less than or equal to the upper limit of normal (ULN)

Ofev PA

Drug Name(s)

Ofev

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND

ii. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary

fibrosis (e.g., radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD) OR

B. BOTH of the following:

i. Patient has a diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD) AND

ii. Patient's diagnosis has been confirmed on high-resolution computed tomography (HRCT) or chest radiography scans OR

C. BOTH of the following:

i. Patient has a diagnosis of chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype AND

ii. Patient's diagnosis has been confirmed on high-resolution computed tomography (HRCT)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of ONE of the following:

A. Idiopathic pulmonary fibrosis (IPF) OR

B. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) OR

C. Chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype AND

3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., pathologist, pulmonologist, radiologist, rheumatologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Oncology Immunotherapy PA – Bavencio

Drug Name(s)

Bavencio

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Updated 10/2024

Oncology Immunotherapy PA – Imfinzi

Drug Name(s)

Imfinzi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Updated 10/2024

Oncology Immunotherapy PA – Jemperli

Drug Name(s)

Jemperli

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Keytruda

Drug Name(s)

Keytruda

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Libtayo

Drug Name(s)

Libtayo

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Opdivo

Drug Name(s)

Opdivo

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Tecentriq

Drug Name(s)

Tecentriq

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Oncology Immunotherapy PA – Yervoy

Drug Name(s)

Yervoy

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. ONE of the following:

a. The requested agent is FDA labeled or supported in CMS approved compendia as first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or CMS approved compendia

supported therapy that are indicated as first-line therapy for the requested agent OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iii. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Opioids ER PA - Buprenorphine Pain

Drug Name(s)

Belbuca

Buprenorphine

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
 - B. Patient has a diagnosis of pain due to sickle cell disease OR
 - C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. ALL of the following:

a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:

1. Diagnosis AND

2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND

b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND

d. ONE of the following:

1. Patient's medication history includes use of an immediate-acting opioid OR

2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR

3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND

e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND

f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months **Other Criteria**:

Opioids ER PA - Fentanyl Patch

Drug Name(s)

Fentanyl

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. ALL of the following:

a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:

1. Diagnosis AND

2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND

b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND

c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND

d. ONE of the following:

1. Patient's medication history includes use of an immediate-acting opioid OR

2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR

3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND

e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months

Opioids ER PA – Hydrocodone

Drug Name(s)

Hydrocodone Bitartrate Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. ALL of the following:

a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:

1. Diagnosis AND

2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND

b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND

c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND

d. ONE of the following:

1. Patient's medication history includes use of an immediate-acting opioid OR

2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR

3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND

e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months

Opioids ER PA – Morphine

Drug Name(s)

Morphine Sulfate Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. ALL of the following:

a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:

1. Diagnosis AND

2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND

b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND

c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND

d. ONE of the following:

1. Patient's medication history includes use of an immediate-acting opioid OR

2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR

3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND

e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months

Opioids ER PA – Oxycodone

Drug Name(s)

Oxycontin

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. ALL of the following:

a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:

1. Diagnosis AND

2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND

b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND

c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND

d. ONE of the following:

1. Patient's medication history includes use of an immediate-acting opioid OR

2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR

3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND

e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Opioids ER PA – Tramadol

Drug Name(s)

Tramadol Hcl Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

iii. ALL of the following:

a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:

1. Diagnosis AND

2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND

b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND

c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND

d. ONE of the following:

1. Patient's medication history includes use of an immediate-acting opioid OR

2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR

3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND

e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Oral Immunotherapy Agents PA – Oralair

Drug Name(s)

Oralair

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. Patient has a diagnosis of allergic rhinitis, with or without conjunctivitis AND
- 2. Patient's diagnosis is confirmed with ONE of the following:
 - A. Positive skin test to ONE of the pollen extracts included in the requested agent OR
 - B. IgE specific antibodies to ONE of the extracts included in the requested agent: Sweet vernal,
 - orchard, perennial rye, Timothy, or Kentucky blue grass AND

3. ONE of the following:

A. Patient has tried and had an inadequate response to TWO standard allergy medications, one of which was an intranasal corticosteroid OR

B. Patient has an intolerance or hypersensitivity to TWO standard allergy medications, one of which was an intranasal corticosteroid OR

C. Patient has an FDA labeled contraindication to TWO standard allergy medications, one of which was an intranasal corticosteroid AND

4. Patient will NOT be using the requested agent in combination with a subcutaneous injectable immunotherapy agent AND

5. The requested agent will be started, or has already been started, 3 to 4 months before the expected onset of the applicable pollen season AND

6. The first dose is given in the clinic/hospital under direct supervision from the provider for a period of at least 30 minutes AND

7. Patient has been prescribed epinephrine auto-injector for at home emergency use

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Standard allergy medications:

Oral antihistamines, oral corticosteroids, intranasal corticosteroids, intranasal antihistamines, or leukotriene inhibitors

Orkambi PA

Drug Name(s)

Orkambi Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND

2. ONE of the following:

A. Patient has the presence of the F508del mutation on both alleles (homozygous) of the CFTR gene confirmed by genetic testing OR

B. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND

3. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of cystic fibrosis AND

3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND 4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for

the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Otezla PA

Drug Name(s)

Otezla

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

1. BOTH of the following:

A. Patient has ONE of the following diagnoses:

- i. Plaque psoriasis OR
- ii. Active psoriatic arthritis AND
- B. ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR

iii. Patient's medication history indicates use of a biologic immunomodulator agent for the same FDA labeled indication OR

iv. Patient has tried and had an inadequate response to at least ONE conventional prerequisite agent for the requested indication OR

v. Patient has an intolerance or hypersensitivity to at least ONE conventional

prerequisite agent for the requested indication OR

vi. Patient has an FDA labeled contraindication to at least ONE conventional prerequisite agent for the requested indication OR

2. Patient has a diagnosis of oral ulcers associated with Behcet's disease (BD)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of plaque psoriasis, active psoriatic arthritis, or oral ulcers associated with Behcet's disease (BD) AND

3. Patient has had clinical benefit with the requested agent (slowing of disease progression or decrease in symptom severity and/or frequency)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Formulary conventional agent required for diagnoses of plaque psoriasis or active psoriatic arthritis

Formulary conventional agents for plaque psoriasis include cyclosporine, methotrexate, tazarotene, topical calcitriol, or topical corticosteroids

Formulary conventional agents for active psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

NO prerequisites are required for a diagnosis of oral ulcers associated with Behcet's disease (BD)

Ozempic PA

Drug Name(s)

Ozempic

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

5. BOTH of the following:

a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND

b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Palynziq PA

Drug Name(s)

Palynziq Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of phenylketonuria (PKU) AND

2. Patient has a baseline blood Phe level greater than 600 micromol/L (10 mg/dL) AND

3. Patient will NOT be using the requested agent in combination with sapropterin for the requested indication AND

4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has a diagnosis of phenylketonuria (PKU) AND
- 3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND

4. Patient will NOT be using the requested agent in combination with sapropterin for the requested indication AND

5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., metabolic or genetic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial approval will be for 9 months, renewal approval will be for 12 months **Other Criteria:**

Panretin PA

Drug Name(s)

Panretin

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of cutaneous lesions associated with AIDS-related Kaposi's sarcoma (KS) OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

- C. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:

a. Patient has a diagnosis of cutaneous lesions associated with AIDSrelated Kaposi's sarcoma (KS) AND

b. Patient does NOT require systemic anti-Kaposi's sarcoma therapy OR

2. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, dermatologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND

iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Pegylated Interferon PA

Drug Name(s)

Pegasys

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic hepatitis B AND BOTH of the following:
 - i. The chronic hepatitis B infection has been confirmed by serological markers AND
 - ii. Patient has NOT been administered the requested agent for more than 48 weeks for
 - the treatment of chronic hepatitis B OR
- B. BOTH of the following:
 - i. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
 - ii. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype OR
- C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

12 months for all other diagnoses. For hep B, hep C see Other Criteria

Other Criteria:

No prior peginterferon alfa use, approve 48 weeks for hepatitis B infection. Prior peginterferon alfa use, approve remainder of 48 weeks of total therapy for hepatitis B infection

Duration of therapy for hepatitis C: Based on FDA approved labeling or AASLD/IDSA guideline supported

Pirfenidone PA Drug Name(s) Pirfenidone Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for initial approval require BOTH of the following: 1. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND

2. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g.,

radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND

3. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., pathologist, pulmonologist, radiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria:**

Posaconazole PA

Drug Name(s)

Noxafil Pak

Posaconazole Inj

Posaconazole Dr

Posaconazole Susp

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of oropharyngeal candidiasis AND ONE of the following:
 - i. Patient has tried and had an inadequate response to fluconazole or an alternative antifungal agent OR

ii. Patient has an intolerance or hypersensitivity to fluconazole or an alternative antifungal agent OR

iii. Patient has an FDA labeled contraindication to fluconazole or an alternative antifungal agent OR

B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR

C. Patient has a diagnosis of invasive Aspergillus AND ONE of the following:

i. Patient has tried and had an inadequate response to an alternative antifungal agent OR

ii. Patient has an intolerance or hypersensitivity to an alternative antifungal agent ORiii. Patient has an FDA labeled contraindication to an alternative antifungal agent OR

D. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

One month for oropharyngeal candidiasis, 6 months for all other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR

B. Patient has a diagnosis of invasive Aspergillus AND patient has continued indicators of active disease (e.g., continued radiologic findings, positive cultures, positive serum galactomannan assay for Aspergillus) OR

C. BOTH of the following:

i. Patient has a diagnosis of oropharyngeal candidiasis AND

ii. Patient has had clinical benefit with the requested agent OR

D. BOTH of the following:

i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Prolia PA

Drug Name(s)

Prolia

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Osteopenia (osteoporosis prophylaxis)

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of:

1. ONE of:

A. Patient's (pt) sex is male or the pt is postmenopausal with a diagnosis of osteoporosis AND BOTH of:

i. Pt's diagnosis was confirmed by ONE of:

- 1. A fragility fracture in the hip or spine OR
- 2. A T-score of -2.5 or lower OR
- 3. A T-score of -1.0 to -2.5 AND ONE of:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR

b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR

c. A FRAX 10-year probability of hip fracture of 3% or greater AND

ii. ONE of:

- 1. Pt is at a very high fracture risk as defined by ONE of:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR
 - c. Pt has had multiple fractures OR

d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR

- e. Pt has a very low T-score (less than -3.0) OR
- f. Pt is at high risk for falls or has a history of injurious falls OR

g. Pt has a very high fracture probability by FRAX (e.g., major

osteoporosis fracture greater than 30%, hip fracture greater than 4.5%)

or by other validated fracture risk algorithm OR

2. ONE of:

a. Pt's medication history includes use of a bisphosphonate OR

b. Pt has an intolerance, FDA labeled contraindication, or

hypersensitivity to a bisphosphonate OR

B. Pt is requesting the agent for osteopenia (osteoporosis prophylaxis) AND ALL of:

i. ONE of:

- 1. Pt's sex is male and the pt is 50 years of age or over OR
- 2. Pt is postmenopausal AND

ii. Pt has a T-score between -1.0 to -2.50 AND

iii. ONE of:

a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR

- b. 10-year probability of a hip fracture 3% and greater per FRAX OR
- c. 10-year probability of a major OP-related fracture 20% and greater per $\ensuremath{\mathsf{FRAX}}$

AND

- iv. ONE of:
 - a. Pt's medication history includes use of a bisphosphonate OR

Criteria continues: See Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR

C. Pt's sex is a female with a diagnosis of breast cancer who is receiving aromatase inhibitor therapy AND ONE of:

i. Pt's medication history includes use of a bisphosphonate OR

ii. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR

D. Pt's sex is male with a diagnosis of prostate cancer receiving androgen deprivation therapy (ADT) AND ONE of:

i. Pt's medication history includes use of a bisphosphonate OR

ii. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR

E. Pt has a diagnosis of glucocorticoid-induced osteoporosis AND ALL of:

i. Pt is either initiating or continuing systemic glucocorticoids in a daily dose equivalent to 7.5 mg or greater of prednisone AND

ii. Pt is expected to remain on glucocorticoids for at least 6 months AND

iii. Pt's diagnosis was confirmed by ONE of:

1. A fragility fracture in the hip or spine OR

2. A T-score of -2.5 or lower OR

3. A T-score of -1.0 to -2.5 AND ONE of the following:

a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR

b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR

c. A FRAX 10-year probability of hip fracture of 3% or greater AND

iv. ONE of:

1. Pt is at a very high fracture risk as defined by ONE of the following:

- a. Pt had a recent fracture (within the past 12 months) OR
- b. Pt had fractures while on FDA approved osteoporosis therapy OR

c. Pt has had multiple fractures OR

d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR

e. Pt has a very low T-score (less than -3.0) OR

f. Pt is at high risk for falls or has a history of injurious falls OR

g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR

, 2. ONE of:

a. Pt's medication history includes use of a bisphosphonate ORb. Pt has an intolerance, FDA labeled contraindication, orhypersensitivity to a bisphosphonate AND

2. ONE of:

A. Pt has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

B. Pt has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent ORC. Prescriber has indicated that the pt is not at risk for hypocalcemia (not including risk associated with the requested agent) AND

3. Pt will NOT be using the requested agent in combination with a bisphosphonate, another form of denosumab (e.g., Xgeva), romosozumab-aqqg, or parathyroid hormone analog (e.g., abaloparatide, teriparatide) for the requested indication AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Promacta PA

Drug Name(s)

Promacta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ONE of the following:

1. Patient (pt) has a diagnosis of persistent or chronic immune (idiopathic) thrombocytopenia (ITP) AND ONE of the following:

A. Pt has tried and had an insufficient response to a corticosteroid or immunoglobulin (IVIg or anti-D) OR

B. Pt has an intolerance or hypersensitivity to a corticosteroid or immunoglobulin (IVIg or anti-D) OR

C. Pt has an FDA labeled contraindication to a corticosteroid or immunoglobulin (IVIg or anti-D) OR

- D. Pt has had an insufficient response to a splenectomy OR
- 2. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND ONE of the following:

A. Pt's platelet count is less than 75 x 10^9/L AND the intent is to increase platelet counts sufficiently to initiate pegylated interferon therapy OR

B. Pt is on concurrent therapy with a pegylated interferon and ribavirin AND is at risk for discontinuing banatitis C therapy due to thrembesytepapia OP

discontinuing hepatitis C therapy due to thrombocytopenia OR

- 3. Pt has a diagnosis of severe aplastic anemia (SAA) AND ALL of the following:
 - A. Pt has at least 2 of the following blood criteria:
 - i. Neutrophils less than 0.5 X 10^9/L OR
 - ii. Platelets less than 30 X 10^9/L OR
 - iii. Reticulocyte count less than 60 X 10^9/L AND
 - B. Pt has at least 1 of the following marrow criteria:

i. Severe hypocellularity is less than 25% OR

ii. Moderate hypocellularity is 25-50% with hematopoietic cells representing less than 30% of residual cells AND

C. ONE of the following:

i. Pt has tried and had an insufficient response to BOTH antithymocyte globulin (ATG) AND cyclosporine therapy OR

ii. BOTH of the following:

1. Pt will be using the requested agent as first-line treatment (i.e., has not been treated with ATG and/or cyclosporine) AND

2. Pt will use the requested agent in combination with standard

immunosuppressive therapy (i.e., ATG AND cyclosporine) OR

4. Pt has another indication that is supported in CMS approved compendia for the requested agent **Age Restriction:**

Prescriber Restrictions:

Coverage Duration:

Initial: 6 months for ITP. Renewal: 12 months for ITP. Other indications, see Other Criteria. **Other Criteria:**

Criteria for renewal approval require BOTH of the following:

1. Pt has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Pt has a diagnosis of persistent or chronic immune (idiopathic) thrombocytopenia (ITP) AND ONE of the following:

i. Pt's platelet count is 50 x 10^9/L or greater OR

ii. Pt's platelet count has increased sufficiently to avoid clinically significant bleeding OR B. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND BOTH of the following:

i. ONE of the following:

1. Pt will be initiating hepatitis C therapy with pegylated interferon and ribavirin OR

2. Pt will be maintaining hepatitis C therapy with pegylated interferon and ribavirin at the same time as the requested agent AND

ii. ONE of the following:

1. Pt's platelet count is 90 x 10^9/L or greater OR

2. Pt's platelet count has increased sufficiently to initiate or maintain pegylated interferon based therapy for the treatment of hepatitis C OR

C. Pt has a diagnosis of severe aplastic anemia (SAA) AND the pt has had clinical benefit with the requested agent OR

D. Pt has another indication that is supported in CMS approved compendia and the pt has had clinical benefit with the requested agent

Initial: 48 weeks for hepatitis C associated thrombocytopenia, 6 months for first-line therapy in severe aplastic anemia, 16 weeks for SAA, 12 months for All other indications

Renewal: 48 weeks for hepatitis C associated thrombocytopenia, 12 months for SAA, 12 months for All other indications

Pulmonary Hypertension PA – Adempas

Drug Name(s)

Adempas

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

- b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
- ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH), WHO Group 4, as determined by a ventilation-perfusion scan and a confirmatory selective pulmonary angiography AND ALL of the following:

i. ONE of the following:

a. Patient is NOT a candidate for surgery OR

b. Patient has had pulmonary endarterectomy AND has persistent or recurrent disease AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg ANDiii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units OR C. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

Initial criteria continues: see Other Criteria Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months **Other Criteria:**

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR c. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Ambrisentan

Drug Name(s)

Ambrisentan

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 (PDE 5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months

Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Bosentan

Drug Name(s)

Bosentan

Tracleer

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Elevated liver enzymes accompanied by signs or symptoms of liver dysfunction/injury or a bilirubin level of 2 times the ULN (upper limit of normal) or greater AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1, as determined by right heart catheterization, AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg ANDiii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 inhibitor (PDE5) plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic

class OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Opsumit

Drug Name(s)

Opsumit

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 (PDE 5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months

Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Sildenafil

Drug Name(s)

Sildenafil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Concurrently taking another phosphodiesterase 5 (PDE 5) inhibitor with the requested agent AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with an endothelin receptor antagonist (ERA) for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy, [except for dual requests for a phosphodiesterase 5 (PDE5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria Age Restriction: Prescriber Restrictions: **Coverage Duration:** Approval will be for 12 months **Other Criteria:**

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy) AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Tadalafil

Drug Name(s)

Alyq

Tadalafil 20Mg Tablet

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Concurrently taking another phosphodiesterase 5 (PDE 5) inhibitor with the requested agent AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with an endothelin receptor antagonist (ERA) for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy, [except for dual requests for a phosphodiesterase 5 (PDE5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy) AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Pulmonary Hypertension PA – Ventavis

Drug Name(s)

Ventavis

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

c. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. Patient is WHO functional class III or IV AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Drug is also subject to Part B versus Part D review.

Prior Authorization Group Description: Pyrimethamine PA Drug Name(s) Pyrimethamine Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require BOTH of the following: 1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND 2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 6 months **Other Criteria:**

Quinine PA

Drug Name(s)

Quinine Sulfate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has ONE of the following diagnoses:

- a. Uncomplicated malaria OR
- b. Babesiosis OR

c. An indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

7 days for malaria, 10 days for babesiosis, 12 months for all other diagnoses **Other Criteria**:

Prior Authorization Group Description: Regranex PA Drug Name(s) Regranex Indications: All Medically-Accepted Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. ONE of the following: a. BOTH of the following: i. Patient has a diagnosis of lower extremity diabetic neuropathic ulcer(s) that extends into the subcutaneous tissue or beyond AND ii. The ulcer(s) intended for treatment has an adequate blood supply OR b. Patient has an indication that is supported in CMS approved compendia for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Repatha PA

Drug Name(s)

Repatha

Repatha Pushtronex System

Repatha Sureclick

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has ONE of the following:

A. A diagnosis of heterozygous familial hypercholesterolemia (HeFH) AND ONE of the following:
 i. Genetic confirmation of one mutant allele at the LDLR, Apo-B, PCSK9, or 1/LDLRAP1
 gene OR

ii. History of LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L) (pretreatment) OR
iii. Patient has clinical manifestations of HeFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthoma, or xanthelasma) OR
iv. Patient has "definite" or "possible" familial hypercholesterolemia as defined by the Simon Broome criteria OR

v. Patient has a Dutch Lipid Clinic Network criteria score of greater than 5 OR vi. Patient has a treated low-density lipoprotein cholesterol (LDL-C) level 100 mg/dL or greater after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy OR

B. A diagnosis of homozygous familial hypercholesterolemia (HoFH) AND ONE of the following:
 i. Genetic confirmation of two mutant alleles at the LDLR, Apo-B, PCSK9, or LDLRAP1
 gene OR

ii. History of untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C 300 mg/dL or greater (7.76 mmol/L or greater) OR

iii. Patient has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma) OR

C. A diagnosis of established cardiovascular disease [angina pectoris, coronary heart disease, myocardial infarction, transient ischemic attacks, cerebrovascular disease (CeVD) or peripheral vascular disease (PVD) or after coronary revascularization or carotid endarterectomy] AND the requested agent will be used to reduce the risk of myocardial infarction, stroke, and coronary revascularization OR

D. A diagnosis of primary hyperlipidemia (not associated with HeFH, HoFH, or established cardiovascular disease) OR

Initial criteria continues: see Other Criteria Age Restriction: Prescriber Restrictions: The agent was prescribed by, or in consultation with, a cardiologist, an endocrinologist, and/or a physician who focuses in the treatment of cardiovascular (CV) risk management and/or lipid disorders

Coverage Duration:

Approval will be for 12 months

Other Criteria:

E. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. Patient has tried and had an inadequate response to a high-intensity statin (i.e., rosuvastatin 20-40 mg or atorvastatin 40-80 mg) OR

- B. Patient has an intolerance to TWO different statins OR
- C. Patient has an FDA labeled contraindication to a statin AND

4. Patient will NOT be using the requested agent in combination with another PCSK9 agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another PCSK9 agent

Prior Authorization Group Description: Ritalin PA Drug Name(s) Methylphenidate Hydrochloride (Ritalin) Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has an FDA labeled indication for the requested agent Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Roflumilast PA

Drug Name(s)

Roflumilast

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

- 1. Patient has an FDA labeled indication for the requested agent AND
- 2. ONE of the following:

A. Patient has tried and had an inadequate response to an agent from two of the following categories:

i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]

ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]

iii. inhaled corticosteroid (ICS) [e.g., fluticasone] OR

B. Patient has an intolerance or hypersensitivity to an agent from two of the following categories:

i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]

ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]

iii. inhaled corticosteroid (ICS) [e.g., fluticasone] OR

C. Patient has an FDA labeled contraindication to an agent from two of the following categories:

i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]

ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]

iii. inhaled corticosteroid (ICS) [e.g., fluticasone]

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Rybelsus PA

Drug Name(s)

Rybelsus

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

- 2. Patient had an ineffective treatment response to an oral diabetes medication
- (e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Samsca PA

Drug Name(s)

Tolvaptan

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the request agent AND Any underlying liver disease, including cirrhosis **Required Medical Information:**

Criteria for approval require ALL of the following:

1. The requested agent was initiated (or re-initiated) in the hospital AND

2. Prior to initiating the requested agent, the patient has or had a diagnosis of clinically significant hypervolemic or euvolemic hyponatremia defined by ONE of the following:

A. Serum sodium is less than 125 mEq/L OR

B. Serum sodium is 125 mEq/L or greater AND patient has symptomatic hyponatremia that has resisted correction with fluid restriction AND

3. Medications known to cause hyponatremia have been evaluated and discontinued when appropriate AND

4. Patient has NOT already received 30 days of therapy with the requested agent following the most recent hospitalization for initiation of therapy AND

5. The requested dose is within the FDA labeled dosing for the requested indication (Recommended starting dose is 15 mg once daily. Dosage may be increased at intervals greater than or equal to 24 hours to 30 mg once daily, and to a maximum of 60 mg once daily as needed to raise serum sodium. Do not administer for more than 30 days to minimize the risk of liver injury.)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 30 days **Other Criteria**:

Sapropterin PA

Drug Name(s)

Javygtor Sapropterin Dihydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of phenylketonuria (PKU) AND

2. Prescriber has submitted a baseline blood Phe level measured prior to initiation of therapy with the requested agent, which is above the recommended levels indicated for the patient's age range or condition AND

3. Patient will NOT be using the requested agent in combination with Palynziq (pegvaliase-pqpz) for the requested indication AND

4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has a diagnosis of phenylketonuria (PKU) AND
- 3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND

4. Patient will NOT be using the requested agent in combination with Palynziq (pegvaliase-pqpz) for the requested indication AND

5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., metabolic or genetic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial: 2 months if dose is 5 to less than 20 mg/kg/day, 1 month if 20 mg/kg/day Renewal: 12 months **Other Criteria**:

Prior Authorization Group Description: Self - Administered Oncology PA Drug Name(s) Abiraterone Acetate Akeega Alecensa Alunbrig Augtyro Ayvakit Balversa Besremi Bexarotene Cap Bosulif Braftovi Brukinsa Cabometyx Calquence Caprelsa Cometriq Copiktra Cotellic Dasatinib Daurismo Erivedge Erleada Erlotinib Hydrochloride Everolimus Exkivity Farydak Fotivda Fruzagla Gavreto Gefitinib Gilotrif Ibrance Iclusig Idhifa Imatinib Mesylate Imbruvica Inlyta Inqovi Inrebic Iwilfin Jakafi

Jaypirca Kisqali Kisqali Femara 200 Dose Kisqali Femara 400 Dose Kisqali Femara 600 Dose Koselugo Krazati Lapatinib Ditosylate Lazcluze Lenalidomide Lenvima 10 Mg Daily Dose Lenvima 12Mg Daily Dose Lenvima 14 Mg Daily Dose Lenvima 18 Mg Daily Dose Lenvima 20 Mg Daily Dose Lenvima 24 Mg Daily Dose Lenvima 4 Mg Daily Dose Lenvima 8 Mg Daily Dose Lonsurf Lorbrena Lumakras Lynparza Lytgobi Matulane Mekinist Mektovi Nerlynx Ninlaro Nubeqa Odomzo Ogsiveo Ojemda Ojjaara Onureg Orgovyx Orserdu Pazopanib Pemazyre Piqray 200Mg Daily Dose Piqray 250Mg Daily Dose Piqray 300Mg Daily Dose Pomalyst Qinlock Retevmo

Revlimid Rezlidhia Rozlytrek Rubraca Rydapt Scemblix Sorafenib Sprycel Stivarga Sunitinib Malate Tabrecta Tafinlar Tagrisso Talzenna Tasigna Tazverik Tepmetko Thalomid Tibsovo Torpenz Tretinoin 10Mg Cap Truqap Tukysa Turalio Vanflyta Venclexta Venclexta Starting Pack Verzenio Vitrakvi Vizimpro Vonjo Voranigo Votrient Welireg Xalkori Xospata Xpovio Xpovio 100 Mg Once Weekly Xpovio 40 Mg Once Weekly Xpovio 40 Mg Twice Weekly Xpovio 60 Mg Once Weekly Xpovio 60 Mg Twice Weekly Xpovio 80 Mg Once Weekly Xpovio 80 Mg Twice Weekly

Xtandi Yonsa Zejula Zelboraf Zolinza Zydelig Zykadia

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND iii. ONE of the following:

a. The requested agent is FDA labeled or supported by compendia as a first-line therapy for the requested indication OR

b. Patient has tried appropriate FDA labeled or compendia supported therapy that are indicated as first-line therapy for the requested indication OR

c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR

d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND

iv. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines AND

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

v. ONE of the following:

a. The requested agent is not Bosulif OR

b. The requested agent is Bosulif AND ONE of the following:

1. Patient's medication history indicates use of imatinib OR Sprycel for the requested indication (if applicable) OR

2. Patient has an intolerance or hypersensitivity to imatinib OR Sprycel OR

3. Patient has an FDA labeled contraindication to imatinib OR Sprycel OR4. CMS approved compendia does not support the use of imatinib ORSprycel for the requested indication OR

5. Prescriber has provided information in support of use of Bosulif over imatinib OR Sprycel for the requested indication AND

- vi. ONE of the following:
 - a. The requested agent is not Calquence OR
 - b. The requested agent is Calquence AND ONE of the following:

1. Patient's medication history indicates use of Brukinsa OR Imbruvica for the requested indication (if applicable) OR

2. Patient has an intolerance or hypersensitivity to Brukinsa OR Imbruvica OR

3. Patient has an FDA labeled contraindication to Brukinsa OR Imbruvica OR

4. CMS approved compendia do not support the use of Brukinsa OR Imbruvica for the requested indication OR

5. Prescriber has provided information in support of use of Calquence over Brukinsa OR Imbruvica for the requested indication

Signifor LAR PA

Drug Name(s)

Signifor Lar

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Severe hepatic impairment (i.e., Child Pugh C)

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of acromegaly AND ONE of the following:

i. Patient had an inadequate response to surgery as indicated by growth hormone and serum IGF-1 levels that are above the reference ranges for the patient's gender and age OR

ii. Patient is NOT a candidate for surgery OR

B. Patient has a diagnosis of Cushing's disease (CD) AND ONE of the following:

- i. Patient had an inadequate response to pituitary surgical resection OR
 - ii. Patient is NOT a candidate for pituitary surgical resection OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. ONE of the following:
 - A. Patient has a diagnosis of acromegaly AND ONE of the following:

i. Patient has growth hormone and serum IGF-1 levels that are within normal limits for patient's gender and age reference range OR

ii. Patient has had clinical improvement (e.g., reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms) OR

B. Patient has a diagnosis of Cushing's disease (CD) AND BOTH of the following:

i. Patient has a urinary free cortisol level less than or equal to the upper limit of normal AND

ii. Patient has had improvement in at least ONE of the following clinical signs and symptoms:

- 1. Fasting plasma glucose OR
- 2. Hemoglobin A1c OR
- 3. Hypertension OR
- 4. Weight OR

C. BOTH of the following:

i. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Initial: Acromegaly - 6 months, CD - 7 months, All other diagnoses - 12 months, Renewal: 12 months **Other Criteria:**

Signifor PA

Drug Name(s)

Signifor

Indications: All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Severe hepatic impairment (i.e., Child Pugh C)

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of Cushing's disease (CD) AND ONE of the following:

i. Patient had an inadequate response to pituitary surgical resection OR

ii. Patient is NOT a candidate for pituitary surgical resection OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of Cushing's disease (CD) AND BOTH of the following:

i. Patient has a urinary free cortisol level less than or equal to the upper limit of normal AND

ii. Patient has had improvement in at least ONE of the following clinical signs and symptoms:

1. Fasting plasma glucose OR

- 2. Hemoglobin A1c OR
- 3. Hypertension OR
- 4. Weight OR

B. BOTH of the following:

i. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Initial approval: 6 months for CD, 12 months for all other diagnoses, Renewal approval: 12 months **Other Criteria:**

Sivextro PA

Drug Name(s)

Sivextro

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following:

A. BOTH of the following:

i. A documented acute bacterial skin and skin structure infection (ABSSSI) defined as a bacterial infection of the skin with a lesion size area of at least 75 cm2 (lesion size measured by the area of redness, edema, or induration) AND

ii. The infection is due to Staphylococcus aureus, Streptococcus pyogenes, Streptococcus agalactiae, Streptococcus anginosus, Streptococcus intermedius, Streptococcus constellatus, or Enterococcus faecalis OR

B. Another indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR

B. The requested agent is NOT prescribed by an infectious disease specialist or the prescriber has NOT consulted with an infectious disease specialist on treatment of this patient AND ONE of the following:

i. There is documentation of resistance to TWO of the following: beta-lactams, macrolides, clindamycin, tetracycline, or co-trimoxazole at the site of infection ORii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams,

macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

- iv. There is documentation of resistance to vancomycin at the site of infection OR
- v. Patient has an intolerance or hypersensitivity to vancomycin OR
- vi. Patient has an FDA labeled contraindication to vancomycin AND

3. Patient will NOT be using the requested agent in combination with linezolid for the same infection AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 days for ABSSSI or 30 days for all other indications

Other Criteria:

Sodium Oxybate PA

Drug Name(s)

Sodium Oxybate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of narcolepsy with cataplexy OR

B. Patient has a diagnosis of narcolepsy with excessive daytime sleepiness AND BOTH of the following:

i. ONE of the following:

- a. Patient is under 18 years of age OR
- b. ONE of the following:

1. Patient has tried and had an inadequate response to modafinil or armodafinil OR

2. Patient has an intolerance or hypersensitivity to modafinil or armodafinil OR

3. Patient has an FDA labeled contraindication to modafinil or armodafinil AND

ii. ONE of the following:

a. Patient has tried and had an inadequate response to ONE standard stimulant agent (e.g., methylphenidate) OR

b. Patient has an intolerance or hypersensitivity to ONE standard stimulant agent (e.g., methylphenidate) OR

c. Patient has an FDA labeled contraindication to ONE standard stimulant agent (e.g., methylphenidate) OR

C. Patient has another indication that is supported in CMS approved compendia for the requested agent

a Postriction:

Age Restriction:

Patient is 7 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Somatostatin Analogs PA – Lanreotide

Drug Name(s)

Somatuline Depot

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

ii. Prescriber states the patient is currently being treated with the requested agent OR

B. ONE of the following:

i. Patient has a diagnosis of acromegaly AND ONE of the following:

a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR

b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR

c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR

ii. Patient has a diagnosis of gastroenteropancreatic neuroendocrine tumors AND BOTH of the following:

- a. The tumors are well or moderately differentiated AND
- b. ONE of the following:
 - 1. The tumors are unresectable, locally advanced OR
 - 2. Patient has metastatic disease OR

iii. Patient has a diagnosis of carcinoid syndrome OR

iv. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. BOTH of the following:

i. ONE of the following:

1. Patient has a diagnosis of acromegaly OR

2. Patient has a diagnosis of metastatic OR unresectable, locally advanced, well or moderately differentiated gastroenteropancreatic neuroendocrine tumors OR

3. Patient has a diagnosis of carcinoid syndrome OR

4. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Somatostatin Analogs PA – Octreotide

Drug Name(s)

Octreotide Acetate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR

B. ONE of the following:

i. Patient has a diagnosis of acromegaly AND ONE of the following:

a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR

b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR

c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR

ii. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR

iii. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR

iv. Patient has a diagnosis of dumping syndrome AND ONE of the following:

a. Patient has tried and had an inadequate response to acarbose OR

- b. Patient has an intolerance or hypersensitivity to acarbose OR
- c. Patient has an FDA labeled contraindication to acarbose OR

v. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal **Other Criteria:**

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of acromegaly OR

B. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR

C. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR

D. Patient has a diagnosis of dumping syndrome OR

E. Patient has another indication that is supported in CMS approved compendia for the requested agent

AND

3. Patient has had clinical benefit with the requested agent AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Somatostatin Analogs PA – Somavert

Drug Name(s)

Somavert

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of acromegaly AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR C. BOTH of the following:

i. ONE of the following:

a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR

b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR

c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by serum IGF-1 levels that are above the reference range AND

ii. ONE of the following:

a. Patient has tried and had an inadequate response to octreotide or Somatuline Depot (lanreotide) OR

b. Patient has an intolerance or hypersensitivity to octreotide or Somatuline Depot (lanreotide) OR

c. Patient has an FDA labeled contraindication to octreotide or Somatuline Depot (lanreotide) AND

2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of acromegaly AND

3. Patient has had clinical benefit with the requested agent AND

4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal **Other Criteria:**

Sovaldi PA

Drug Name(s)

Sovaldi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND

2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND

3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND

4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND

5. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR

D. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration:**

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria**:

Spravato PA

Drug Name(s)

Spravato 56Mg Dose

Spravato 84Mg Dose

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. BOTH of the following:

a. ONE of the following:

i. Patient has a diagnosis of treatment-resistant depression (TRD) OR

ii. Patient has a diagnosis of major depressive disorder (MDD) with acute suicidal ideation or behavior AND

b. The requested agent will be used in combination with an oral antidepressant AND

2. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent OR

- c. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:

a. Patient has a diagnosis of treatment-resistant depression (TRD) AND

b. Patient has tried and had an inadequate response to at least two

different oral antidepressants (e.g., SSRIs, SNRIs) OR

2. Patient has a diagnosis of major depressive disorder (MDD) with acute suicidal ideation or behavior AND

ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., psychiatrist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND

iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Strensiq PA

Drug Name(s)

Strensiq Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

A. Perinatal or infantile-onset hypophosphatasia OR

B. Juvenile-onset hypophosphatasia AND

2. Patient has documentation (i.e., medical records) of clinical manifestations to support the diagnosis of hypophosphatasia at the age of onset prior to age 18 (e.g., vitamin B6-dependent seizures, skeletal abnormalities such as rachitic chest deformity leading to respiratory problems or bowed arms/legs, "failure to thrive") AND

3. Patient has documentation (i.e., medical records) of radiographic imaging to support the diagnosis of hypophosphatasia at the age of onset prior to age 18 (e.g., infantile rickets, alveolar bone loss, craniosynostosis, fractures) AND

4. Patient has documentation (i.e., medical records) of confirmed mutation(s) in the ALPL gene that encodes the tissue non-specific isoenzyme of alkaline phosphatase (TNSALP) AND

5. Patient has documentation (i.e., medical records) of a measured total serum alkaline phosphatase (ALP) level that is below the normal lab reference range for age and sex AND

6. Patient has documentation (i.e., medical records) of ONE of the following:

A. Elevated urine concentration of phosphoethanolamine (PEA) OR

B. Elevated serum concentration of pyridoxal 5'-phosphate (PLP) in the absence of vitamin supplements within one week prior to the test OR

C. Elevated urinary inorganic pyrophosphate (PPi) AND

7. The requested dose is within FDA labeled dosing (based on the patient's weight) for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist or geneticist with expertise in metabolic bone diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has ONE of the following diagnoses:

A. Perinatal or infantile-onset hypophosphatasia OR

B. Juvenile-onset hypophosphatasia AND

3. There is documentation (i.e., medical records) that the patient has had a decrease from baseline (before treatment with the requested agent) in at least ONE of the following levels:

a. Urine concentration of phosphoethanolamine (PEA) OR

b. Serum concentration of pyridoxal 5'-phosphate (PLP) in the absence of vitamin supplements within one week prior to the test OR

c. Urinary inorganic pyrophosphate (PPi) AND

4. Patient has documentation (i.e., medical records) of clinical improvement and/or stabilization with the requested agent (e.g., improvement in respiratory status, growth, pain, radiographic findings, other symptoms associated with the disease) AND

5. The requested dose is within FDA labeled dosing (based on the patient's weight) for the requested indication

Substrate Reduction Therapy PA – Miglustat

Drug Name(s)

Miglustat

Yargesa Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:

A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR

B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND

2. Prescriber has drawn baseline measurements of hemoglobin level, platelet count, liver volume, and spleen volume AND

3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:

A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR

- B. Thrombocytopenia (defined as platelet count of less than 100,000 per microliter) OR
- C. Hepatomegaly OR
- D. Splenomegaly OR
- E. Growth failure (i.e., growth velocity is below the standard mean for age) OR
- F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND

3. Patient has had improvements or stabilization with the requested agent as indicated by ONE of the following:

- A. Spleen volume OR
- B. Hemoglobin level OR
- C. Liver volume OR
- D. Platelet count OR
- E. Growth OR
- F. Bone pain or crisis

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration: Approval will be for 12 months **Other Criteria:**

Tasimelteon Capsule PA

Drug Name(s)

Tasimelteon

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. Patient has a diagnosis of Non-24-hour sleep-wake disorder AND

ii. Patient is totally blind (i.e., no light perception) OR

B. BOTH of the following:

i. Patient has a diagnosis of Smith-Magenis Syndrome (SMS) confirmed by the presence of ONE of the following genetic mutations:

- A. A heterozygous deletion of 17p11.2 OR
- B. A heterozygous pathogenic variant involving RAI1 AND

ii. The requested agent is being used to treat nighttime sleep disturbances associated with SMS

Age Restriction:

For diagnosis of Non-24-hour sleep-wake disorder, patient is 18 years of age or over. For diagnosis of Smith-Magenis Syndrome (SMS), patient is 16 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist, sleep specialist, psychiatrist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration**:

Approval will be for 12 months **Other Criteria**:

Teriparatide PA

Drug Name(s)

Forteo

Teriparatide

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

- A. Postmenopausal with osteoporosis OR
- B. Patient's sex is male with primary or hypogonadal osteoporosis OR
- C. Osteoporosis with sustained systemic glucocorticoid therapy AND
- 2. Patient's diagnosis was confirmed by ONE of the following:
 - A. A fragility fracture in the hip or spine OR
 - B. A T-score of -2.5 or lower OR

C. A T-score of -1.0 to -2.5 AND ONE of the following:

- i. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
- ii. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
- iii. A FRAX 10-year probability of hip fracture of 3% or greater AND
- 3. ONE of the following:
 - A. Patient is at a very high fracture risk as defined by ONE of the following:
 - i. Patient had a recent fracture (within the past 12 months) OR
 - ii. Patient had fractures while on FDA approved osteoporosis therapy OR
 - iii. Patient has had multiple fractures OR

iv. Patient had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR

v. Patient has a very low T-score (less than -3.0) OR

vi. Patient is at high risk for falls or has a history of injurious falls OR

vii. Patient has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture

risk algorithm OR

B. ONE of the following:

i. Patient has tried and had an inadequate response to a bisphosphonate OR

- ii. Patient has an intolerance or hypersensitivity to a bisphosphonate OR
- iii. Patient has an FDA labeled contraindication to a bisphosphonate AND

4. Patient will NOT be using the requested agent in combination with a bisphosphonate, denosumab (e.g., Prolia, Xgeva), romosozumab-aqqg, or another parathyroid hormone analog (e.g., abaloparatide) for the requested indication AND

Criteria continues: see Other Criteria Age Restriction:

Prescriber Restrictions:

Coverage Duration:

No prior teriparatide and/or Tymlos use approve 2 years, Prior use - see Other Criteria **Other Criteria**:

5. The requested dose is within FDA labeled dosing for the requested indication AND

6. ONE of the following:

A. Patient has never received treatment with teriparatide or Tymlos (abaloparatide) ORB. Patient has been previously treated with teriparatide or Tymlos (abaloparatide) AND ONE of the following:

i. The total cumulative duration of treatment with teriparatide and Tymlos (abaloparatide) has NOT exceeded 2 years OR

ii. Patient has received 2 years or more of treatment with teriparatide, and remains at or has returned to having a high risk for fracture

Prior teriparatide and/or Tymlos use approve remainder of 2 years of total cumulative therapy. Approve 1 year if patient has received 2 years or more teriparatide

Tetrabenazine PA

Drug Name(s)

Tetrabenazine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

- A. Patient has a diagnosis of chorea associated with Huntington's disease OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
- 2. ONE of the following:
 - A. Patient does NOT have a current diagnosis of depression OR
 - B. Patient has a current diagnosis of depression and is being treated for depression AND
- 3. ONE of the following:
 - A. Patient does NOT have a diagnosis of suicidal ideation and/or behavior OR
 - B. Patient has a diagnosis of suicidal ideation and/or behavior and must NOT be actively suicidal AND

4. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) AND

5. Patient will NOT be using the requested agent in combination with reserpine

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Tobramycin neb PA

Drug Name(s)

Tobramycin Neb

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND

2. Documentation has been provided that indicates the patient has a Pseudomonas aeruginosa respiratory infection AND

3. ONE of the following:

a. Patient is NOT currently (within the past 60 days) being treated with another inhaled antibiotic (e.g., inhaled aztreonam, inhaled tobramycin) OR

b. Patient is currently (within the past 60 days) being treated with another inhaled antibiotic (e.g., inhaled aztreonam, inhaled tobramycin) AND ONE of the following:

i. Prescriber has confirmed that the other inhaled antibiotic will be discontinued, and that therapy will be continued only with the requested agent OR

ii. Prescriber has provided information in support of another inhaled antibiotic therapy used concurrently with or alternating with (i.e., continuous alternating therapy) the requested agent

Drug is also subject to Part B versus Part D review. Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Topical Diclofenac 3% Gel PA Drug Name(s) Diclofenac Sodium 3% Gel Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has a diagnosis of actinic keratosis (AK) Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 3 months **Other Criteria:**

Topical Retinoids PA – Tazarotene

Drug Name(s)

Tazarotene

Tazorac
Indications:
Indications:
All Medically-Accepted Indications.
Off-Label Uses:
Exclusion Criteria:
Requested agent will be used for cosmetic purposes
Required Medical Information:
Criteria for approval require the following:
1. ONE of the following:
a. Patient has an FDA labeled indication for the requested agent OR
b. Patient has an indication that is supported in CMS approved compendia for the requested agent
Age Restriction:
Prescriber Restrictions:
Coverage Duration:

Approval will be for 12 months

Topical Retinoids PA – Tretinoin

Drug Name(s)

Avita

Tretinoin Cream, Gel

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for cosmetic purposes

Required Medical Information:

Criteria for approval require the following:

- 1. ONE of the following:
 - a. Patient has an FDA labeled indication for the requested agent OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Trelstar PA

Drug Name(s)

Trelstar Mixject

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR C. BOTH of the following:

i. Patient is NOT currently being treated with the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND

3. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction: Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Trientine PA

Drug Name(s)

Trientine Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of Wilson's disease confirmed by ONE of the following:

- A. Confirmation of genetic mutation of the ATP7B gene OR
- B. Patient has TWO or more of the following:
 - i. Presence of hepatic abnormality (e.g., acute liver failure, cirrhosis, fatty liver)
 - ii. Presence of Kayser-Fleischer rings

iii. Serum ceruloplasmin level less than 20 mg/dL

- iv. Basal urinary copper excretion greater than 40 mcg/24 hours or the testing laboratory's upper limit of normal
- v. Hepatic parenchymal copper content greater than 40 mcg/g dry weight
- vi. Presence of neurological symptoms (e.g., dystonia, hypertonia, rigidity with tremors,
- dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) AND
- 2. ONE of the following:
 - A. Patient has tried and had an inadequate response to penicillamine OR
 - B. Patient has an intolerance or hypersensitivity to penicillamine OR
 - C. Patient has an FDA labeled contraindication to penicillamine

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

- 2. Patient has a diagnosis of Wilson's disease AND
- 3. Patient has had clinical benefit with the requested agent as evidenced by ONE of the following:
 - A. Improvement and/or stabilization in hepatic abnormality OR
 - B. Reduction in Kayser-Fleischer rings OR
 - C. Improvement and/or stabilization in neurological symptoms (e.g., dystonia, hypertonia,

rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) OR D. Basal urinary copper excretion greater than 200 mcg/24 hours

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist, neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Trikafta PA

Drug Name(s)

Trikafta Indications: All FDA-Approved Indications. Off-Label Uses: Exclusion Criteria: Required Medical Information: Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of cystic fibrosis AND 2. ONE of the following:

A. Patient has the presence of the F508del mutation in at least ONE allele (heterozygous OR homozygous) of the CFTR gene confirmed by genetic testing OR

B. Patient has ONE of the CFTR gene mutations or a mutation in the CFTR gene that is responsive based on in vitro data, as indicated in the FDA label, confirmed by genetic testing OR

C. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND

3. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of cystic fibrosis AND

3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND

4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria:**

Trulicity PA

Drug Name(s)

Trulicity

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

5. BOTH of the following:

a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND

b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Tymlos PA

Drug Name(s)

Tymlos

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient (pt) has ONE of the following:

A. Postmenopausal with osteoporosis OR

- B. Pt's sex is male with osteoporosis AND
- 2. BOTH of the following:

A. Pt's diagnosis was confirmed by ONE of the following:

i. A fragility fracture in the hip or spine OR

ii. A T-score of -2.5 or lower OR

iii. A T-score of -1.0 to -2.5 AND ONE of the following:

a. A fragility fracture of proximal humerus, pelvis, or distal forearm OR

b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR

c. A FRAX 10-year probability of hip fracture of 3% or greater AND

B. ONE of the following:

i. Pt is at a very high fracture risk as defined by ONE of the following:

a. Pt had a recent fracture (within the past 12 months) OR

b. Pt had fractures while on FDA approved osteoporosis therapy OR

c. Pt has had multiple fractures OR

d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR

e. Pt has a very low T-score (less than -3.0) OR

f. Pt is at high risk for falls or has a history of injurious falls OR

g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis

fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR

ii. ONE of the following:

a. Pt has tried and had an inadequate response to a bisphosphonate OR

b. Pt has an intolerance or hypersensitivity to a bisphosphonate OR

c. Pt has an FDA labeled contraindication to a bisphosphonate AND

3. Pt will NOT be using the requested agent in combination with a bisphosphonate, denosumab (e.g., Prolia, Xgeva), romosozumab-aqqg, or another parathyroid hormone analog (e.g., teriparatide) for the requested indication AND

4. The requested dose is within FDA labeled dosing for the requested indication AND

5. The total cumulative duration of treatment with teriparatide and Tymlos (abaloparatide) has not exceeded 2 years

Age Restriction: Prescriber Restrictions: Coverage Duration: No prior Tymlos and/or teriparatide use approve 2 years, Prior use - see Other Criteria Other Criteria: Prior Tymlos and/or teriparatide use approve remainder of 2 years of total cumulative therapy

Ubrelvy PA

Drug Name(s)

Ubrelvy

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

- 1. Patient has a diagnosis of migraine AND
- 2. The requested agent is being used for the treatment of acute migraine with or without aura AND
- 3. ONE of the following:

A. Patient has tried and had an inadequate response to a triptan (e.g., sumatriptan, rizatriptan) agent OR

- B. Patient has an intolerance, or hypersensitivity to a triptan OR
- C. Patient has an FDA labeled contraindication to a triptan AND

4. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of migraine AND

- 3. The requested agent is being used for the treatment of acute migraine with or without aura AND
- 4. Patient has had clinical benefit with the requested agent AND

5. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP)

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Urea Cycle Disorders PA - Sodium Phenylbutyrate

Drug Name(s)

Sodium Phenylbutyrate

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of ONE of the following:

a. Urea cycle disorder with neonatal-onset involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase OR

b. Urea cycle disorder with late-onset and history of hyperammonemic encephalopathy involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase AND

2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., geneticist, metabolic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Prior Authorization Group Description: Viberzi PA Drug Name(s) Viberzi Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has a diagnosis of irritable bowel syndrome with diarrhea (IBS-D) Age Restriction: **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months **Other Criteria:**

Victoza PA

Drug Name(s)

Victoza

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. BOTH of the following:

i. ONE of the following:

1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR

2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR

3. Patient has an intolerance or hypersensitivity to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

4. Patient has an FDA labeled contraindication to an oral diabetes medication

(e.g., metformin, an agent containing metformin, glipizide) OR

5. BOTH of the following:

a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND

b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Prescriber Restrictions: Coverage Duration: Approval will be for 12 months Other Criteria:

Voriconazole PA

Drug Name(s)

Voriconazole

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of invasive Aspergillus OR
- B. Patient has a serious infection caused by Scedosporium apiospermum or Fusarium species OR
- C. Patient has a diagnosis of esophageal candidiasis or candidemia in nonneutropenic patient AND ONE of the following:
 - i. Patient has tried and had an inadequate response to fluconazole or an alternative antifungal agent OR
 - ii. Patient has an intolerance or hypersensitivity to fluconazole or an alternative antifungal agent OR
 - iii. Patient has an FDA labeled contraindication to fluconazole or an alternative antifungal agent OR
- D. Patient has a diagnosis of blastomycosis AND ONE of the following:
 - i. Patient has tried and had an inadequate response to itraconazole OR
 - ii. Patient has an intolerance or hypersensitivity to itraconazole OR
 - iii. Patient has an FDA labeled contraindication to itraconazole OR
- E. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- F. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

One month for esophageal candidiasis, 6 months for all other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of invasive Aspergillus, a serious infection caused by Scedosporium apiospermum or Fusarium species, esophageal candidiasis, candidemia in nonneutropenic patient, or blastomycosis and patient has continued indicators of active disease (e.g., continued radiologic findings, positive cultures, positive serum galactomannan assay for Aspergillus) OR B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR

C. BOTH of the following:

i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Vosevi PA

Drug Name(s)

Vosevi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND

2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND

3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND

4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND

5. If genotype 1, the patient's subtype has been identified and provided

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis **Coverage Duration**:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria:**

Vyndamax PA

Drug Name(s)

Vyndamax

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND

2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND

3. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND

3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND

4. Patient has had clinical benefit with the requested agent AND

5. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Vyndaqel PA

Drug Name(s)

Vyndaqel

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND

2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND

3. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND

3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND

4. Patient has had clinical benefit with the requested agent AND

5. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months **Other Criteria**:

Prior Authorization Group Description: Xermelo PA Drug Name(s) Xermelo Indications: All FDA-Approved Indications. Off-Label Uses: **Exclusion Criteria: Required Medical Information:** Criteria for initial approval require ALL of the following: 1. Patient has a diagnosis of carcinoid syndrome diarrhea AND 2. Patient has tried and had an inadequate response to treatment with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide]) AND 3. The requested agent will be used in combination with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide]) Criteria for renewal approval require ALL of the following: 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of carcinoid syndrome diarrhea AND

3. Patient has had clinical benefit with the requested agent (e.g., reduction in the average number of daily bowel movements) AND

4. The requested agent will be used in combination with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide])

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Xgeva PA

Drug Name(s)

Xgeva

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

- 1. ONE of the following:
 - A. Patient has a diagnosis of multiple myeloma AND BOTH of the following:
 - i. The requested agent will be used for the prevention of skeletal-related events AND
 - ii. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

B. Patient has a diagnosis of prostate cancer AND ALL of the following:

- i. The requested agent will be used for the prevention of skeletal-related events AND
- ii. Patient has bone metastases AND

iii. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

C. Patient has a solid tumor cancer diagnosis (e.g., thyroid, non-small cell lung, kidney cancer, or breast cancer) AND ALL of the following:

i. The requested agent will be used for the prevention of skeletal-related events AND

ii. Patient has bone metastases AND

iii. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

D. Patient has a diagnosis of giant cell tumor of bone AND ONE of the following:

i. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

ii. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

iii. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

E. Patient has a diagnosis of hypercalcemia of malignancy AND

2. Patient will NOT be using the requested agent in combination with Prolia (denosumab) AND

3. The requested dose is within FDA labeled dosing for the requested indication

Prior Authorization Group Description: Xifaxan PA Drug Name(s) Xifaxan Indications: All FDA-Approved Indications. **Off-Label Uses: Exclusion Criteria:** FDA labeled contraindications to the requested agent **Required Medical Information:** Criteria for approval require the following: 1. Patient has ONE of the following: a. A diagnosis of irritable bowel syndrome with diarrhea (IBS-D) OR b. A diagnosis of hepatic encephalopathy [reduction in risk of overt hepatic encephalopathy (HE) recurrence] OR c. BOTH of the following: i. A diagnosis of traveler's diarrhea (TD) AND ii. The traveler's diarrhea is caused by noninvasive strains of Escherichia coli Age Restriction: For diagnosis of traveler's diarrhea (TD), patient is within the FDA labeled age for the requested agent **Prescriber Restrictions: Coverage Duration:** Approval will be for 12 months

Xolair PA

Drug Name(s)

Xolair

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. ONE of the following:

- A. Patient has a diagnosis of moderate to severe persistent asthma AND ALL of the following:
 - i. ONE of the following:
 - a. Patient is 6 to less than 12 years of age AND BOTH of the following:
 - I. Patient's pretreatment IgE level is 30 IU/mL to 1300 IU/mL AND
 - II. Patient's weight is 20 kg to 150 kg OR
 - b. Patient is 12 years of age or over AND BOTH of the following:
 - I. Patient's pretreatment IgE level is 30 IU/mL to 700 IU/mL AND
 - II. Patient's weight is 30 kg to 150 kg AND

iii. Allergic asthma has been confirmed by a positive skin test or in vitro reactivity test (RAST) to a perennial aeroallergen AND

iv. ONE of the following:

a. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an asthma control therapy OR

B. Patient has a diagnosis of chronic idiopathic urticaria AND BOTH of the following:

i. Patient has had over 6 weeks of hives and itching AND

ii. ONE of the following:

a. Patient has tried and had an inadequate response to maximum tolerable H1 antihistamine therapy OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to H1 antihistamine therapy OR

C. Patient has a diagnosis of nasal polyps AND BOTH of the following:

i. ONE of the following:

a. Patient has tried and had an inadequate response to an intranasal corticosteroid OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an intranasal corticosteroid AND

ii. ONE of the following:

a. The requested agent will be used in combination with an intranasal corticosteroid OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an intranasal corticosteroid OR

Initial criteria continues: see Other Criteria

Age Restriction:

For diagnosis of moderate to severe persistent asthma, patient is 6 years of age or over. For diagnosis of chronic idiopathic urticaria, patient is 12 years of age or over. For diagnosis of nasal polyps, patient is 18 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist, otolaryngologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

D. Patient has a diagnosis of IgE-mediated food allergy AND ALL of the following:

i. Patient is using the requested agent for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods AND

ii. IgE-mediated food allergy has been confirmed by an allergy diagnostic test (e.g., skin prick test, serum specific IgE test, oral food challenge) AND

iii. Patient will avoid known food allergens while treated with the requested agent AND

iv. The requested agent will NOT be used for the emergency treatment of allergic reactions, including anaphylaxis AND

 Patient will NOT be using the requested agent in combination with Dupixent or an injectable Interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
 The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of moderate to severe persistent asthma AND ALL of the following:

i. Patient's weight is within the FDA indicated range for their age (i.e., 20 kg to 150 kg for patients age 6 to less than 12 years and 30 kg to 150 kg for patients 12 years of age or over) AND

ii. Patient has had clinical benefit with the requested agent AND

iii. ONE of the following:

a. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an asthma control therapy OR

B. Patient has a diagnosis of chronic idiopathic urticaria AND the following:

- a. Patient has had clinical benefit with the requested agent OR
- C. Patient has a diagnosis of nasal polyps AND the following:
 - a. Patient has had clinical benefit with the requested agent OR
- D. Patient has a diagnosis of IgE-mediated food allergy AND ALL of the following:

a. Patient is using the requested agent for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods AND

b. Patient has had clinical benefit with the requested agent AND

c. Patient will avoid known food allergens while treated with the requested agent AND

d. The requested agent will NOT be used for the emergency treatment of allergic reactions, including anaphylaxis AND

3. Patient will NOT be using the requested agent in combination with Dupixent or an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND 4. The requested dose is within FDA labeled dosing for the requested indication

Zepatier PA

Drug Name(s)

Zepatier

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND

2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND

3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND

4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND

5. If genotype 1, the patient's subtype has been identified and provided AND

6. If genotype 1a, the prescriber has tested the patient for the presence of virus with NS5A resistanceassociated polymorphisms AND

7. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR

C. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR

D. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported **Other Criteria:**

Zokinvy PA

Drug Name(s)

Zokinvy

Indications:

All FDA-Approved Indications.

Off-Label Uses:

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. Patient has a diagnosis of Hutchinson-Gilford progeria syndrome (HGPS) AND

ii. Genetic testing has confirmed a pathogenic variant in the LMNA gene that results in production of progerin OR

B. Patient has a diagnosis of processing-deficient progeroid laminopathy AND ONE of the following:

i. Genetic testing has confirmed heterozygous LMNA mutation with progerin-like protein accumulation OR

ii. Genetic testing has confirmed homozygous or compound heterozygous ZMPSTE24 mutations

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of ONE of the following:

A. Hutchinson-Gilford progeria syndrome (HGPS) OR

B. Processing-deficient progeroid laminopathies with either: heterozygous LMNA mutation with progerin-like protein accumulation OR homozygous or compound heterozygous ZMPSTE24 mutations AND

3. Patient has had clinical benefit with the requested agent

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist, geneticist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months